

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

02

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>A61K 38/17, 39/395, G01N 33/577, 33/68</b>		<b>A1</b>	(11) International Publication Number: <b>WO 97/03687</b>
			(43) International Publication Date: <b>6 February 1997 (06.02.97)</b>
(21) International Application Number: <b>PCT/US96/12010</b>		(81) Designated States: <b>AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>	
(22) International Filing Date: <b>19 July 1996 (19.07.96)</b>		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: <b>08/505,606      21 July 1995 (21.07.95)      US</b>			
(60) Parent Application or Grant (63) Related by Continuation <b>US      08/505,606 (CIP)</b> <b>Filed on      21 July 1995 (21.07.95)</b>			
(71) Applicant (for all designated States except US): <b>BIODIN, INC.</b> <b>[US/US]; 14 Cambridge Center, Cambridge, MA 02142 (US).</b>			
(72) Inventors; and (75) Inventors/Applicants (for US only): <b>BROWNING, Jeffrey, L.</b> <b>[US/US]; 32 Milton Road, Brookline, MA 02146 (US).</b> <b>BENJAMIN, Christopher, D. [US/US]; 2 Oak Hill Lane,</b> <b>Beverly, MA 01915 (US). HOCHMAN, Paula, S. [US/US];</b> <b>95 Collidge Street, Brookline, MA 02146 (US).</b>			
(74) Agent: <b>FLYNN, Kerry, A.; Bion, Inc., 14 Cambridge Center,</b> <b>Cambridge, MA 02142 (US).</b>			
(54) Title: <b>SOLUBLE LYMPHOTOXIN-<math>\beta</math> RECEPTORS AND ANTI-LYMPHOTOXIN RECEPTOR AND LIGAND ANTIBODIES, AS THERAPEUTIC AGENTS FOR THE TREATMENT OF IMMUNOLOGICAL DISEASE</b>			
<b>1      SQPQAVPPYA SENQTCRDQE KEYYEPQHRI CCSRCPPGTY VSAKCSRIRD      50</b> <b>51      TVCATCAENS YNEHWNYLTI QCLCRPCDPV MGLEEIAPCT SKRKTQCRCQ      100</b> <b>101      PGMFCAAWAL ECTHCELLSD CPPGTEAELK DEVGKGNHNC VPCKAGHFQN      150</b> <b>151      TSSPSARCQP HTRCENQGLV EAAPGTAQSD TTCKNPLEPL PPEMSGT      197</b>			
(57) Abstract			
<p>This invention relates to compositions and methods comprising "lymphotoxin-<math>\beta</math> receptor blocking agents", which block lymphotoxin-<math>\beta</math> receptor signalling. Lymphotoxin-<math>\beta</math> receptor blocking agents are useful for treating lymphocyte-mediated immunological diseases, and more particularly, for inhibiting Th1 cell-mediated immune responses. This invention relates to soluble forms of the lymphotoxin-<math>\beta</math> receptor extracellular domain that act as lymphotoxin-<math>\beta</math> receptor blocking agents. This invention also relates to the use of antibodies directed against either the lymphotoxin-<math>\beta</math> receptor or its ligand, surface lymphotoxin, that act as lymphotoxin-<math>\beta</math> receptor blocking agents. A novel screening method for selecting soluble receptors, antibodies and other agents that block LT-<math>\beta</math> receptor signalling is provided.</p>			

BEST AVAILABLE COPY

66

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

SOLUBLE LYMPHOTOXIN- $\beta$  RECEPTORS AND  
ANTI-LYMPHOTOXIN RECEPTOR AND LIGAND ANTIBODIES AS THERAPEUTIC  
AGENTS FOR  
THE TREATMENT OF IMMUNOLOGICAL DISEASE

TECHNICAL FIELD OF THE INVENTION

This invention relates to compositions and methods comprising "lymphotoxin- $\beta$  receptor blocking agents", which block lymphotoxin- $\beta$  receptor signalling. Lymphotoxin- $\beta$  receptor blocking agents are useful for treating lymphocyte-mediated immunological diseases, and more particularly, for inhibiting Th1 cell-mediated immune responses. This invention relates to soluble forms of the lymphotoxin- $\beta$  receptor extracellular domain that act as lymphotoxin- $\beta$  receptor blocking agents. This  
10 invention also relates to the use of antibodies directed against either the lymphotoxin- $\beta$  receptor or its ligand, surface lymphotoxin, that act as lymphotoxin- $\beta$  receptor blocking agents. A novel screening method for selecting soluble receptors, antibodies and other agents that block LT- $\beta$  receptor signalling  
15 is provided.

BACKGROUND OF THE INVENTION

The pattern of cytokines released at the onset of an immune challenge can affect the subsequent choice of which immune effector pathways are activated. The choice between  
20 immune effector mechanisms is mediated by CD4-positive helper T lymphocytes (T helper cells or Th cells). Th cells interact with antigen-presenting cells (APCs), which display peptide fragments of processed foreign antigen in association with MHC class II molecules on their surfaces. Th cells are activated  
25 when they recognize particular epitopes of a foreign antigen displayed on the appropriate APC surface for which the Th cells express a specific receptor. Activated Th cells, in turn,

- 2 -

secrete cytokines (lymphokines) which activate appropriate immune effector mechanisms.

Th cells can activate diverse effector mechanisms, including killer T cell activation B cell antibody production  
5 and macrophage activation. The choice between effector mechanisms is mediated largely by which cytokines are produced by the activated Th cells.

Th cells can be divided into three subgroups based on their cytokine secretion patterns (Fitch et al., Ann. Rev. Immunol., 11, pp. 29-48 (1993)). These subgroups are called  
10 Th0, Th1 and Th2. In the mouse, non-stimulated "naive" T helper cells produce IL-2. Short term stimulation leads to Th0 precursor cells, which produce a wide range of cytokines including IFN- $\gamma$ , IL-2, IL-4, IL-5 and IL-10. Chronically-  
15 stimulated Th0 cells can differentiate into either Th1 or Th2 cell types, whereupon the cytokine expression pattern changes.

Some cytokines are released by both Th1 and Th2 cells (e.g., IL-3, GM-CSF and TNF). Other cytokines are made exclusively by one or the other Th cell subgroup. The  
20 specialized effects of T helper cell subgroups were first recognized in mouse. A similar subdivision of T helper cells also exists in humans (Romagnani et al., Ann. Rev. Immunol., 12, pp. 227-57 (1994)).

Th1 cells produce LT- $\alpha$ , IL-2 and IFN- $\gamma$ . In humans,  
25 the Th1 pattern of cytokine secretion has been generally associated with cellular immunity and resistance to infection. The Th1 cytokines tend to activate macrophages and certain inflammatory responses such as Type IV "delayed type" hypersensitivity (see below). Th1 cytokines play an important  
30 role in cellular rejection of tissue grafts and organ transplants.

Th2 cells produce the cytokines IL-4, IL-5, IL-6 and IL-10. Th2 cytokines increase eosinophil and mast cell production and promote the full expansion and maturation of B  
35 cells (Howard et al., "T cell-derived cytokines and their receptors", Fundamental Immunology, 3d ed., Raven Press, New York (1993)). Th2 cytokines also enhance antibody production, including IgE antibodies associated with allergic responses and

- 3 -

anti-graft antibodies. Th2 cells may also participate in immune suppression and tolerance to persistent antigens.

Th1- and Th2-associated cytokines play a role in certain hypersensitivity responses -- inappropriate or  
5 disproportionate immune responses evoked upon contact with a previously encountered antigen. There are four recognized types of hypersensitivity (Roitt et al., Immunology, pp. 19.1-22.12 (Mosby-Year Book Europe Ltd., 3d ed. 1993)).

Type I "immediate hypersensitivity" involves allergen-  
10 induced Th2 cell activation and Th2 cytokine release. The Th2 cytokine IL-4 stimulates B cells to undergo isotype switching to produce IgE, which activates mast cells to produce acute inflammatory reactions such as those which lead to eczema, asthma and rhinitis.

15 Types II and III hypersensitivity are caused by IgG and IgM antibodies directed against cell surface or specific tissue antigens (Type II) or soluble serum antigens (Type III). These types of hypersensitivity reactions are not thought to be mediated by Th cells.

20 Type IV "delayed type" hypersensitivity (DTH) is Th1 cell mediated. DTH reactions take more than 12 hours to develop and are referred to as "cell-mediated" because they can be transferred between mice by transferring Th1 cells but not serum alone. Type IV DTH responses are generally classified into  
25 three types: contact, tuberculin-type and granulomatous hypersensitivity.

Many cell-mediated responses that can cause disease are inducible in healthy mice by transferring lymphocytes from a diseased mouse (e.g., insulin-dependent diabetes and  
30 experimental autoimmune encephalitis). This feature distinguishes Type IV DTH from the other three types of hypersensitivity, which are humoral immune responses caused primarily by antibodies which can be transferred in cell-free serum.

35 T helper cells also participate in the regulation of de novo immunoglobulin isotype switching. Different Th subsets may influence the relative proportion of immunoglobulins of a given isotype produced in response to immune challenge. For

- 4 -

example, the Th2 cytokine IL-4 can switch activated B cells to the IgG1 isotype and suppress other isotypes. As discussed above, IL-4 also activates IgE overproduction in type I hypersensitivity reactions. The Th2 cytokine IL-5 induces the IgA isotype. These Th2 cytokine effects on isotype switching are counter-balanced by IFN- $\gamma$  produced by Th1 cells.

The differential patterns of cytokines secreted by Th1 and Th2 cells appear to direct a response towards different immune effector mechanisms. The switch that activates either a cell-mediated or humoral effector mechanism is sensitized by cross-suppression between Th1 and Th2 cells: IFN- $\gamma$  produced by Th1 cells inhibits Th2 cell proliferation and Th2 cell-secreted IL-10 appears to reduce cytokine secretion from Th1 cells.

Depending on the relative affinities of the cytokines for their molecular targets, the Th1 and Th2 negative regulatory circuits may amplify the effects of small concentration differences between Th1 and Th2 cytokines. An amplified Th1 or Th2 cytokine signal may trigger the switch between cell-mediated or humoral effector mechanisms based on small changes in the relative concentrations of Th1 and Th2 cytokines. The ability to control this switch by modulating the relative concentrations of Th1 and Th2 cytokines would be useful for treating imbalances in a variety of Th1 and Th2 cell-dependent immune responses which can lead to immune disorders and diseases.

Pathological Th1 responses are associated with a number of organ-specific and systemic autoimmune conditions, chronic inflammatory diseases, and delayed type hypersensitivity reactions. As discussed above, Th1 responses also contribute to cellular responses leading to grafted tissue and transplanted organ rejection.

The treatment of these various Th1 cell-based immunological conditions to date has generally employed immunomodulatory and immuno suppressive agents as well as a number of drugs with poorly characterized mechanisms (e.g., gold or penicillamine). Three general immunosuppressive agents used currently are steroids, cyclosporine and azathioprine.

Steroids are pleiotropic anti-inflammatory agents which suppress activated macrophages and inhibit the activity of

- 5 -

antigen presenting cells in ways which reverse many of the effects of the Th1 cytokine IFN- $\gamma$ . Cyclosporine -- a potent immunosuppressive agent -- suppresses cytokine production and reduces the expression of IL-2 receptors on lymphocytes during  
5 their activation. Azathioprine is an anti-proliferative agent which inhibits DNA synthesis. These non-specific immunosuppressive agents are generally required in high doses which increase their toxicity (e.g. nephro- and hepatotoxicity) and cause adverse side effects. They are thus unsuitable for  
10 long term therapies.

To address the problems caused by conventional treatments with non-specific immunosuppressive agents, many current therapeutic strategies aim at suppressing or activating selective aspects of the immune system. An especially  
15 attractive goal is the manipulation of the balance between Th1 and Th2 cytokines to shift the balance between cell-mediated and humoral effector mechanisms.

To accomplish a shift between cell-mediated and humoral effector mechanisms, it would be useful to be able to  
20 modulate the activity of a molecule that can shift the relative activities of Th1 and Th2 cell subclasses. Candidates for such molecules include the cytokines and their receptors. Recent data suggest that LT- $\alpha$ , IL-12, IFN- $\alpha$  and IFN- $\gamma$  favor the development of Th1 responses, whereas IL-1 and IL-4 polarize a  
25 response towards a Th2 effector mechanism (Romagnani et al., Ann. Rev. Immunol., 12 , pp. 227-57 (1994)).

Many of the Th cell cytokines are pleiotropic regulators of immune development and function, and inhibiting their production would have deleterious effects on non-T cell  
30 mediated responses. A desirable and effective target for selectively modulating the choice between Th1 and Th2 effector mechanisms has not been identified.

#### SUMMARY OF THE INVENTION

The present invention solves the problems referred to  
35 above by providing pharmaceutical compositions and methods for treating immunological diseases by inhibiting lymphotoxin- $\beta$  receptor (LT- $\beta$ -R) signalling using lymphotoxin- $\beta$  receptor

- 6 -

blocking agents. More particularly, the compositions and methods comprising LT- $\beta$ -R blocking agents are useful for inhibiting Th1 cell-mediated immune responses such as, for example, inflammatory bowel syndrome.

5 In one embodiment, soluble forms of the lymphotoxin- $\beta$  receptor extracellular domain that act as LT- $\beta$ -R blocking agents are provided. The preferred compositions and methods of this embodiment comprise a recombinant lymphotoxin- $\beta$  receptor fusion protein that has the LT- $\beta$ -R extracellular ligand binding domain  
10 fused to an immunoglobulin constant heavy chain domain. More preferably, the LT- $\beta$ -R ligand binding domain is fused to a human IgG Fc domain.

In another embodiment of this invention, antibodies that act as LT- $\beta$ -R blocking agents are provided. Preferred  
15 compositions and methods of this embodiment comprise one or more antibodies directed against the lymphotoxin- $\beta$  receptor. More preferably, the antibody is a monoclonal antibody. Other preferred compositions and methods of this embodiment comprise one or more antibodies directed against surface lymphotoxin.  
20 More preferably, the antibody is a monoclonal antibody directed against lymphotoxin- $\beta$ .

This invention further provides a novel screening process for selecting LT- $\beta$ -R blocking agents -- such as soluble forms of the LT- $\beta$ -R, anti-LT Abs and anti-LT- $\beta$ -R Abs. This  
25 screening process involves performing tumor cell cytotoxicity assays that monitor LT- $\beta$ -R signalling. The assay makes use of the increased sensitivity of human adenocarcinoma cells to ligand- or antibody-induced LT- $\beta$ -R signalling in the presence of an LT- $\beta$ -R activating agent (such as LT- $\alpha$ 1/ $\beta$ 2) in a tumor  
30 cytotoxicity assay.

LT- $\beta$ -R blocking agents inhibit the cytotoxic effects of LT- $\alpha$ / $\beta$  heteromeric complexes (or other LT- $\beta$ -R activating agents) on tumor cells. The procedure used to test putative  
LT- $\beta$ -R blocking agents is exemplified for the case of anti-LT- $\beta$ -  
35 R antibodies (in the presence of the LT- $\beta$ -R activating agents LT- $\alpha$ 1/ $\beta$ 2) and comprises the following steps:

1) Tumor cells (e.g., HT29 human adenocarcinoma cells) are cultured for several days in media containing IFN- $\gamma$



- 7 -

and purified LT- $\alpha$ 1/ $\beta$ 2 in the presence or absence of the particular anti-LT- $\beta$ -R Ab being assayed;

2) The cells are treated with a dye that stains living cells; and

5 3) The number of stained cells is quantitated to determine the fraction of tumor cells killed in the presence of LT- $\alpha$ 1/ $\beta$ 2, IFN- $\gamma$  and the test anti-LT- $\beta$ -R Ab in each sample. Alternatively, the number of surviving cells can be determined by any of a number of well-known assays which measure cell  
10 viability, such as  $^3\text{H}$ -thymidine incorporation into DNA. An anti-LT- $\beta$ -R Ab (or an Ab combination) that decreases the percentage of tumor cells killed in this assay by at least 20% is a LT- $\beta$ -R blocking agent within the scope of this invention.

This cytolytic assay may be performed using LT- $\alpha$ / $\beta$   
15 heteromeric complexes and other LT- $\beta$ -R activating agents, either alone or in combination. The assay can also be adapted as required to identify new LT- $\beta$ -R blocking agents.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**Figure 1.** The sequence of the extracellular portion of the  
20 human LT- $\beta$  receptor which encodes the ligand binding domain.

**Figure 2.** A soluble murine LT- $\beta$  receptor coupled to the human IgG1 Fc domain (mLT- $\beta$ -R-Fc) blocks LT- $\beta$ -R signalling in mouse WEHI 164 cells induced by soluble murine LT- $\alpha$ / $\beta$  ligand. WEHI 164 cells are killed as a function of increasing LT ligand  
25 (mLT- $\alpha$ / $\beta$ ) concentration. Soluble mLT- $\beta$ -R-Fc (10  $\mu\text{g}/\text{ml}$ ) blocks this LT ligand-induced cell death. A soluble murine TNF receptor fusion protein (p55TNF-R-Fc) has little effect on blocking LT- $\alpha$ / $\beta$ -activated cell death. Growth was quantitated after three days by measuring the optical density (OD 550) of  
30 reacted MTT, which is proportional to cell number.

**Figure 3.** An antibody directed against human LT- $\beta$ -R (BDA8 mAb) blocks the interaction between soluble LT ligand and LT- $\beta$ -R on a human cell surface. The growth of WiDr tumor cells is blocked by a combination of IFN- $\gamma$  and soluble LT- $\alpha$ 1/ $\beta$ 2 ligand. The anti-LT- $\beta$ -R antibody BDA8 blocks the ability  
35 of LT- $\alpha$ 1/ $\beta$ 2 ligand to inhibit the growth of WiDr tumor cells.

- 8 -

Solid symbols show cell growth in the presence of IgG1 control mAb (10  $\mu$ g/ml). Open symbols show the effects of anti-LT- $\beta$ -R mAb BDA8 (10  $\mu$ g/ml).

**Figure 4.** An antibody directed against human LT- $\beta$  (B9 mAb) blocks the interaction between cell surface LT- $\alpha/\beta$  ligand and soluble LT- $\beta$  receptor (hLT- $\beta$ -R-Fc; 2 $\mu$ g/ml). Surface bound LT- $\beta$ -R-Fc was detected using phycoerythrin-labelled donkey anti-human IgG and FACS analysis. The mean fluorescence intensity of the resultant peak is plotted as channel number. Dotted line shows the mean fluorescence intensity corresponding to the amount of receptor bound in the absence of the B9 mAb.

**Figure 5.** The effects of a LT- $\beta$ -R blocking agent (mLT- $\beta$ -R-Fc) on ear swelling in a mouse contact delayed type hypersensitivity model (DTH). The graph shows the increase in ear thickness measured 24 hours following 0.2% DNFB antigen challenge onto the ears of sensitized mice. Each symbol represents a separate experiment. All experiments utilized 7-8 animals per point except those demarcated with a diamond, which used only 4 animals per point. Mice treated with buffer (PBS) and with 20 mg/kg of a control IgG fusion protein (LFA3-Fc) served as negative controls. Mice treated with 8 mg/kg of an anti-VLA4 mAb (PS/2 mAb), which inhibits contact DTH ear swelling, served as positive controls.

**Figure 6** is a graph of the weight change observed in mice 14 days after treatment with mLT- $\beta$ R-lg and hLFA3-lg fusion proteins.

**Figure 7** is a graph of the change in colon length observed in mice 14 days after treatment with mLT- $\beta$ R-lg and hLFA3-lg fusion proteins.

**Figure 8** is a time course of the body weight of mice following injection of CD45RB<sup>low</sup>CD4 positive T cells; CD45RB<sup>high</sup>CD4 positive T cells; CD45RB<sup>high</sup> and in LT $\beta$ R-lg; and CD45RB<sup>high</sup> and hLFA3-lg.

**Figure 9** is a graphical representation of the mean and standard deviations of the body weights observed following the treatments in Figures 8-11.

- 9 -

**Figure 10** is a representation of the increase in footpad thickness of mice injected with negative and positive controls and mLT $\beta$ R-Ig.

#### DETAILED DESCRIPTION OF THE INVENTION

5           In order that the invention herein described may be fully understood, the following detailed description is set forth.

          The term "cytokine" refers to a molecule which mediates interactions between cells. A "lymphokine" is a  
10 cytokine released by lymphocytes.

          The term "T helper (Th) cells" refers to a functional subclass of T cells which help to generate cytotoxic T cells and which cooperate with B cells to stimulate antibody production. Helper T cells recognize antigen in association with class II  
15 MHC molecules.

          The term "Th1" refers to a subclass of T helper cells that produce LT- $\alpha$ , interferon- $\gamma$  and IL-2 (and other cytokines) and which elicit inflammatory reactions associated with a cellular, i.e. non-immunoglobulin, response to a challenge.

20           The term "Th2" refers to a subclass of T helper cells that produce cytokines, including IL-4, IL-5, IL-6 and IL-10, which are associated with an immunoglobulin (humoral) response to an immune challenge.

          The term "cell mediated" refers to those immunological  
25 events that result from the direct effects of T cells and their products to produce a response. This type of response is generally (but not exclusively) associated with the Th1 class of T cells. Not included in this category would be the helper effects of T cells on B cell differentiation and B cell  
30 expansion, which are generally associated with the Th2 class of T cells.

          The term "delayed type hypersensitivity (DTH)" refers to an immunological response that is characterized by a slow response to an antigen with the full effect manifesting itself  
35 over a 1-3 day period. This slow response is in contrast to the relatively fast response seen in an immunoglobulin-mediated (humoral) allergic reaction. There are three types of DTH

- 10 -

reactions: contact hypersensitivity, tuberculin-type hypersensitivity and granulomatous reactions.

The terms "immunoglobulin response" or "humoral response" refer to the immunological response of an animal to a foreign antigen whereby the animal produces antibodies to the foreign antigen. The Th2 class of T helper cells are critical to the efficient production of high affinity antibodies.

The term "Fc domain" of an antibody refers to a part of the molecule comprising the hinge, CH2 and CH3 domains, but lacking the antigen binding sites. The term is also meant to include the equivalent regions of an IgM or other antibody isotype.

The term "anti-LT- $\beta$  receptor antibody" refers to any antibody that specifically binds to at least one epitope of the LT- $\beta$  receptor.

The term "anti-LT antibody" refers to any antibody that specifically binds to at least one epitope of LT- $\alpha$ , LT- $\beta$  or a LT- $\alpha/\beta$  complex.

The term "LT- $\beta$ -R signalling" refers to molecular reactions associated with the LT- $\beta$ -R pathway and subsequent molecular reactions which result therefrom.

The term "LT- $\beta$ -R blocking agent" refers to an agent that can diminish ligand binding to LT- $\beta$ -R, cell surface LT- $\beta$ -R clustering or LT- $\beta$ -R signalling, or that can influence how the LT- $\beta$ -R signal is interpreted within the cell.

A LT- $\beta$ -R blocking agent that acts at the step of ligand-receptor binding can inhibit LT ligand binding to the LT- $\beta$ -R by at least 20%. A LT- $\beta$ -R blocking agent that acts after the step of ligand-receptor binding can inhibit the cytotoxic effects of LT- $\beta$ -R activation on a tumor cell by at least 20%. Examples of LT- $\beta$ -R blocking agents include soluble LT- $\beta$ -R-Fc molecules, and anti-LT- $\alpha$ , anti-LT- $\beta$ , anti-LT- $\alpha/\beta$  and anti-LT- $\beta$ -R Abs. Preferably, the antibodies do not cross-react with the secreted form of LT- $\alpha$ .

The term "LT- $\beta$ -R biological activity" refers to: 1) the ability of the LT- $\beta$ -R molecule or derivative to compete for soluble or surface LT ligand binding with soluble or surface LT- $\beta$ -R molecules; or

- 11 -

2) the ability to stimulate an immune regulatory response or cytotoxic activity in common with a native LT- $\beta$ -R molecule.

The terms "LT- $\alpha/\beta$  heteromeric complex" and "LT heteromeric complex" refer to a stable association between at least one LT- $\alpha$  and one or more LT- $\beta$  subunits, including soluble, mutant, altered and chimeric forms of one or more of the subunits. The subunits can associate through electrostatic, van der Waals, or covalent interactions. Preferably, the LT- $\alpha/\beta$  heteromeric complex has at least two adjacent LT- $\beta$  subunits and lacks adjacent LT- $\alpha$  subunits. When the LT- $\alpha/\beta$  heteromeric complex serves as a LT- $\beta$ -R activating agent in a cell growth assay, the complex is preferably soluble and has the stoichiometry LT- $\alpha_1/\beta_2$ . Soluble LT- $\alpha/\beta$  heteromeric complexes lack a transmembrane domain and can be secreted by an appropriate host cell which has been engineered to express LT- $\alpha$  and/or LT- $\beta$  subunits (Crowe et al., J. Immunol. Methods, 168, pp. 79-89 (1994)).

The term "LT ligand" refers to a LT heteromeric complex or derivative thereof that can specifically bind to the LT- $\beta$  receptor.

The term "LT- $\beta$ -R ligand binding domain" refers to the portion or portions of the LT- $\beta$ -R that are involved in specific recognition of and interaction with a LT ligand.

The terms "surface LT- $\alpha/\beta$  complex" and "surface LT complex" refer to a complex comprising LT- $\alpha$  and membrane-bound LT- $\beta$  subunits -- including mutant, altered and chimeric forms of one or more of the subunits -- which is displayed on the cell surface. "Surface LT ligand" refers to a surface LT complex or derivative thereof that can specifically bind to the LT- $\beta$  receptor.

The term "subject" refers to an animal, or to one or more cells derived from an animal. Preferably, the animal is a mammal. Cells may be in any form, including but not limited to cells retained in tissue, cell clusters, immortalized, transfected or transformed cells, and cells derived from an animal that have been physically or phenotypically altered.

**Lymphotoxin- $\beta$ : A Member of the TNF Family**

- 12 -

Tumor Necrosis Factor (TNF)-related cytokines have emerged as a large family of pleiotropic mediators of host defense and immune regulation. Members of this family exist in membrane-bound forms which act locally through cell-cell contact, or as secreted proteins which can act on distant targets. A parallel family of TNF-related receptors react with these cytokines and trigger a variety of pathways including cell death, cell proliferation, tissue differentiation and proinflammatory responses.

10           TNF, lymphotoxin- $\alpha$  (LT- $\alpha$ , also called TNF- $\beta$ ) and lymphotoxin- $\beta$  (LT- $\beta$ ) are members of the TNF family of ligands, which also includes the ligands to the Fas, CD27, CD30, CD40, OX-40 and 4-1BB receptors (Smith et al., Cell, 76, pp. 959-62 (1994)). Signalling by several members of the TNF family --  
15 including TNF, LT- $\alpha$ , LT- $\beta$  and Fas -- can induce tumor cell death by necrosis or apoptosis (programmed cell death). In non-tumorigenic cells, TNF and many of the TNF family ligand-receptor interactions influence immune system development and responses to various  
20 immune challenges.

Most TNF family ligands are found as a membrane-bound form on the cell surface. TNF and LT- $\alpha$  are found in both secreted and membrane-associated surface forms in humans. Surface TNF has a transmembrane region that is proteolytically  
25 cleaved to generate the secreted form. In contrast, surface LT- $\alpha$  lacks a transmembrane region. Membrane-associated LT- $\alpha$  is tethered to the cell surface as a heteromeric complex with LT- $\beta$ , a related transmembrane polypeptide, in a LT- $\alpha$ / $\beta$  complex.

Most membrane-associated LT- $\alpha$ / $\beta$  complexes ("surface  
30 LT") have a LT- $\alpha$ 1/ $\beta$ 2 stoichiometry (Browning et al., Cell, 72, pp. 847-56 (1993); Browning et al., J. Immunol., 154, pp. 33-46 (1995)). Surface LT ligands do not bind TNF-R with high affinity and do not activate TNF-R signalling. Another TNF-related receptor, called the LT- $\beta$  receptor (LT- $\beta$ -R), binds these  
35 surface lymphotoxin complexes with high affinity (Crowe et al., Science, 264, pp. 707-10 (1994)).

LT- $\beta$ -R signalling, like TNF-R signalling, has anti-proliferative effects and can be cytotoxic to tumor cells. In

- 13 -

applicants' co-pending United States application serial number 08/378,968, compositions and methods for selectively stimulating LT- $\beta$ -R using LT- $\beta$ -R activating agents are disclosed. LT- $\beta$ -R activating agents are useful for inhibiting tumor cell growth without co-activating TNF-R-induced proinflammatory or immunoregulatory pathways.

In non-tumor cells, TNF and TNF-related cytokines are active in a wide variety of immune responses. Both TNF and LT- $\alpha$  ligands bind to and activate TNF receptors (p55 or p60 and p75 or p80; herein called "TNF-R"). TNF and LT- $\alpha$  are produced by macrophages in an early and rapid response to microbial infection which enhances the microbicidal activity of macrophages and neutrophils. TNF and LT- $\alpha$  made by macrophages or cytotoxic T lymphocytes (CTLs or "killer T cells") bind to TNF receptors on target cell surfaces and trigger the death of susceptible cells.

TNF and TNF-related cytokines can also initiate inflammatory cascades in response to infection or stress. The release of TNF, LT- $\alpha$  and IFN- $\gamma$  changes the adhesion properties between the vascular endothelial cells and certain lymphocyte types. Increased adhesion facilitates phagocyte and leucocyte migration from the bloodstream into the tissues surrounding an inflammation site. Similar inflammatory reactions play a major role in cellular rejection of tissue grafts and organ transplants, and in certain immune disorders.

Cell surface lymphotoxin (LT) complexes have been characterized in CD4<sup>+</sup> T cell hybridoma cells (II-23.D7) which express high levels of LT (Browning et al., J. Immunol., 147, pp. 1230-37 (1991); Androlewicz et al., J. Biol. Chem., 267, pp. 2542-47 (1992)). The expression and biological roles of LT- $\beta$ -R, LT subunits and surface LT complexes have been reviewed in C.F. Ware et al., "The ligands and receptors of the lymphotoxin system", in Pathways for Cytolysis, Current Topics Microbiol. Immunol., Springer-Verlag, pp. 175-218 (1995).

LT- $\alpha$  expression is induced and LT- $\alpha$  secreted primarily by activated T and B lymphocytes and natural killer (NK) cells. Among the T helper cell subclasses, LT- $\alpha$  appears to be produced by Th1 but not Th2 cells. LT- $\alpha$  has also been detected in

- 14 -

melanocytes. Microglia and T cells in lesions of multiple sclerosis patients can also be stained with anti-LT- $\alpha$  antisera.

Lymphotoxin- $\beta$  (also called p33), has been identified on the surface of T lymphocytes, T cell lines, B cell lines and lymphokine-activated killer (LAK) cells. LT- $\beta$  is the subject of applicants' co-pending international applications PCT/US91/04588, published January 9, 1992 as WO 92/00329; and PCT/US93/11669, published June 23, 1994 as WO 94/13808, which are herein incorporated by reference.

Surface LT complexes are primarily expressed by activated T and B lymphocytes and natural killer (NK) cells as defined by FACS analysis or immunohistology using anti-LT- $\alpha$  antibodies or soluble LT- $\beta$ -R-Fc fusion proteins. Surface LT has also been described on human cytotoxic T lymphocyte (CTL) clones, activated peripheral mononuclear lymphocytes (PML), IL-2-activated peripheral blood lymphocytes (LAK cells), pokeweed mitogen-activated or anti-CD40- activated peripheral B lymphocytes (PBL) and various lymphoid tumors of T and B cell lineage. Engagement of alloantigen-bearing target cells specifically induces surface LT expression by CD8<sup>+</sup> and CD4<sup>+</sup> CTL clones.

The LT- $\beta$  receptor, a member of the TNF family of receptors, specifically binds to surface LT ligands. LT- $\beta$ -R binds LT heteromeric complexes (predominantly LT- $\alpha$ 1/ $\beta$ 2 and LT- $\alpha$ 2/ $\beta$ 1) but does not bind TNF or LT- $\alpha$  (Crowe et al., Science, 264, pp. 707-10 (1994)). Signalling by LT- $\beta$ -R may play a role in peripheral lymphoid organ development and in humoral immune responses.

Studies on LT- $\beta$ -R expression are in their early stages. LT- $\beta$ -R mRNAs are found in human spleen, thymus and other major organs. LT- $\beta$ -R expression patterns are similar to those reported for p55-TNF-R except that LT- $\beta$ -R is lacking on peripheral blood T cells and T cell lines.

#### **Production of soluble LT complexes**

Soluble LT- $\alpha$ / $\beta$  heteromeric complexes comprise LT- $\alpha$  subunits which have been changed from a membrane-bound to a soluble form. These complexes are described in detail in



- 15 -

applicants' co-pending international application  
(PCT/US93/11669, published June 23, 1994 as WO 94/13808).

Soluble LT- $\alpha$  peptides are defined by the amino acid sequence of  
lymphotoxin- $\alpha$  wherein the sequence is cleaved at any point

- 5 between the end of the transmembrane region (i.e. at about amino  
acid #44) and the first TNF homology region (i.e. at amino acid  
#88) according to the numbering system of Browning et al., Cell,  
72, pp. 847-56 (1993).

- Soluble LT- $\alpha$  polypeptides may be produced by
- 10 truncating the N-terminus of LT- $\alpha$  to remove the cytoplasmic tail  
and transmembrane region (Crowe et al., Science, 264, pp. 707-  
710 (1994)). Alternatively, the transmembrane domain may be  
inactivated by deletion, or by substitution of the normally  
hydrophobic amino acid residues which comprise a transmembrane  
15 domain with hydrophilic ones. In either case, a substantially  
hydrophilic hydropathy profile is created which will reduce  
lipid affinity and improve aqueous solubility. Deletion of the  
transmembrane domain is preferred over substitution with  
hydrophilic amino acid residues because it avoids introducing  
20 potentially immunogenic epitopes.

- The deleted or inactivated transmembrane domain may be  
replaced with or attached to a type I leader sequence (e.g. the  
VCAM-1 leader) such that the protein is secreted beginning with  
a sequence anywhere from between val40 to pro88. Soluble LT- $\alpha$   
25 polypeptides may include any number of well-known leader  
sequences at the N-terminus. Such a sequence would allow the  
peptides to be expressed and targeted to the secretion pathway  
in a eukaryotic system. See, e.g., Ernst et al., United States  
Patent No. 5,082,783 (1992).

- 30 Soluble LT- $\alpha$ / $\beta$  heteromeric complexes may be produced  
by co-transfecting a suitable host cell with DNA encoding LT- $\alpha$   
and soluble LT- $\beta$  (Crowe et al., J. Immunol. Methods, 168, pp.  
79-89 (1994)). Soluble LT- $\beta$  secreted in the absence of LT- $\alpha$  is  
highly oligomerized. However, when co-expressed with LT- $\alpha$ , a 70  
35 kDa trimeric-like structure is formed which contains both  
proteins. It is also possible to produce soluble LT- $\alpha$ 1/ $\alpha$ 2  
heteromeric complexes by transfecting a cell line which normally

- 16 -

expresses only LT- $\alpha$  (such as the RPMI 1788 cells discussed above) with a gene encoding a soluble LT- $\beta$  polypeptide.

LT- $\alpha$  and LT- $\beta$  polypeptides may be separately synthesized, denatured using mild detergents, mixed together and renatured by removing the detergent to form mixed LT heteromeric complexes which can be separated (see below).

#### Purification of LT- $\alpha$ 1/ $\beta$ 2 Complexes

Soluble LT- $\alpha$ 1/ $\beta$ 2 heteromeric complexes are separated from co-expression complexes comprising a different subunit stoichiometry by chromatography using TNF and LT- $\beta$  receptors as affinity purification reagents. The TNF receptors only bind within  $\alpha/\alpha$  clefts of LT complexes. The LT- $\beta$  receptor binds with high affinity to  $\beta/\beta$  clefts, and with lower affinity to  $\alpha/\beta$  clefts of heteromeric LT- $\alpha$ / $\beta$  complexes. Accordingly, LT- $\alpha$ 3 and LT- $\alpha$ 2/ $\beta$ 1 will bind to TNF-R. The LT- $\beta$ -R can also bind LT- $\alpha$ 2/ $\beta$ 1 trimers (within the  $\alpha/\beta$  clefts) but cannot bind LT- $\alpha$ 3. In addition, the LT- $\beta$ -R (but not TNF-R) binds LT- $\alpha$ 1/ $\beta$ 2 and LT- $\beta$ n (the exact composition of such preparation is unknown, however, they are large aggregates).

The receptor affinity reagents can be prepared as either a soluble extracellular domain (see for example Loetscher et al., J. Biol. Chem., 266, pp. 18324-29 (1991)), or as chimeric proteins with the extracellular ligand binding domain coupled to an immunoglobulin Fc domain (Loetscher et al., J. Biol. Chem., 266, pp. 18324-29 (1991); Crowe et al., Science, 264, pp. 707-710 (1994)). Receptors are coupled to affinity matrices by chemical cross-linking using routine procedures.

There are two schemes by which the LT- $\alpha$ 1/ $\beta$ 2 ligand can be purified using receptors and immuno-affinity chromatography. In the first scheme, a supernatant from an appropriate expression system co-expressing both LT- $\alpha$  and the truncated LT- $\beta$  form is passed over a TNF-R column. The TNF-R will bind LT- $\alpha$ 3 and LT- $\alpha$ 2/ $\beta$ 1 trimers. The flow through from the TNF-R column will contain LT- $\beta$ (n) and LT- $\alpha$ 1/ $\beta$ 2.

In the second scheme, all LT- $\beta$ -containing forms (LT- $\beta$ (n), LT- $\alpha$ 1/ $\beta$ 2 and LT- $\alpha$ 2/ $\beta$ 1) are bound to and eluted from a LT- $\beta$ -R column using classical methods such as chaotrope or pH

- 17 -

change. (LT- $\alpha$ 3 flows through this column). The eluate is neutralized or the chaotrope removed, and the eluate is then passed over a TNF-R column, which binds only to the LT- $\alpha$ 2/ $\beta$ 1 trimers. The flow through of this column will contain LT- $\alpha$ (n) and LT- $\alpha$ 1/ $\beta$ 2 trimers.

In both cases, pure LT- $\alpha$ 1/ $\beta$ 2 trimers can be separated from LT- $\alpha$  by subsequent gel filtration and/or ion exchange chromatographic procedures known to the art.

Alternatively, different forms of LT- $\alpha$ / $\beta$  heteromeric complexes can be separated and purified by a variety of conventional chromatographic means. It may also be preferable to combine a series of conventional purification schemes with one of the immunoaffinity purification steps described above.

#### Screening For LT- $\beta$ -R Blocking Agents

In one embodiment of this invention, the LT- $\beta$ -R blocking agent comprises an antibody (Ab) directed against LT- $\beta$ -R that inhibits LT- $\beta$ -R signalling. Preferably, the anti-LT- $\beta$ -R Ab is a monoclonal antibody (mAb). One such inhibitory anti-LT- $\beta$ -R mAb is BDA8 mAb.

Inhibitory anti-LT- $\beta$ -R Abs and other LT- $\beta$ -R blocking agents can be identified using screening methods that detect the ability of one or more agents either to bind to the LT- $\beta$ -R or LT ligand, or to inhibit the effects of LT- $\beta$ -R signalling on cells.

One screening method makes use of the cytotoxic effects of LT- $\beta$ -R signalling on tumor cells bearing the LT- $\beta$ -R. Tumor cells are exposed to one or more LT- $\beta$ -R activating agents to induce LT- $\beta$ -R signalling. LT- $\beta$ -R activating agents include LT- $\alpha$ / $\beta$  heteromeric complexes (preferably soluble LT- $\alpha$ 1/ $\beta$ 2) in the presence of IFN- $\gamma$ , or an activating anti-LT- $\beta$ -R Ab (see below; also described in applicants' co-pending United States application serial number 08/378,968). Antibodies and other agents that can block LT- $\beta$ -R signalling are selected based on their ability to inhibit the cytotoxic effect of LT- $\beta$ -R signalling on tumor cells in the following assay:

1) Tumor cells such as HT29 cells are cultured for three to four days in a series of tissue culture wells

- 18 -

containing media and at least one LT- $\beta$ -R activating agent in the presence or absence of serial dilutions of the agent being tested;

2) A vital dye stain which measures mitochondrial  
5 function such as MTT is added to the tumor cell mixture and reacted for several hours;

3) The optical density of the mixture in each well is quantitated at 550 nm wavelength light (OD 550). The OD 550 is proportional to the number of tumor cells remaining in the  
10 presence of the LT- $\beta$ -R activating agent and the test LT- $\beta$ -R blocking agent in each well. An agent or combination of agents that can reduce LT- $\beta$ -R-activated tumor cell cytotoxicity by at least 20% in this assay is a LT- $\beta$ -R blocking agent within the scope of this invention.

15 Any agent or combination of agents that activate LT- $\beta$ -R signalling can be used in the above assay to identify LT- $\beta$ -R blocking agents. LT- $\beta$ -R activating agents that induce LT- $\beta$ -R signalling (such as activating anti-LT- $\beta$ -R mAbs) can be selected based on their ability -- alone or in combination with other  
20 agents -- to potentiate tumor cell cytotoxicity using the tumor cell assay described above.

Another method for selecting an LT- $\beta$ -R blocking agent is to monitor the ability of the putative agent to directly interfere with LT ligand-receptor binding. An agent or  
25 combination of agents that can block ligand-receptor binding by at least 20% is an LT- $\beta$ -R blocking agent within the scope of this invention.

Any of a number of assays that measure the strength of ligand-receptor binding can be used to perform competition  
30 assays with putative LT- $\beta$ -R blocking agents. The strength of the binding between a receptor and ligand can be measured using an enzyme-linked immunoadsorption assay (ELISA) or a radio-immunoassay (RIA). Specific binding may also be measured by fluorescently labelling antibody-antigen complexes and  
35 performing fluorescence-activated cell sorting (FACS) analysis, or by performing other such immunodetection methods, all of which are techniques well known in the art.

- 19 -

The ligand-receptor binding interaction may also be measured with the BIAcore instrument (Pharmacia Biosensor) which exploits plasmon resonance detection (Zhou et al., Biochemistry, 32, pp. 8193-98 (1993); Faegerstram and O'Shannessy, "Surface plasmon resonance detection in affinity technologies", in Handbook of Affinity Chromatography, pp. 229-52, Marcel Dekker, Inc., New York (1993)).

The BIAcore technology allows one to bind receptor to a gold surface and to flow ligand over it. Plasmon resonance detection gives direct quantitation of the amount of mass bound to the surface in real time. This technique yields both on and off rate constants and thus a ligand-receptor dissociation constant and affinity constant can be directly determined in the presence and absence of the putative LT- $\beta$ -R blocking agent.

With any of these or other techniques for measuring receptor-ligand interactions, one can evaluate the ability of a LT- $\beta$ -R blocking agent, alone or in combination with other agents, to inhibit binding of surface or soluble LT ligands to surface or soluble LT- $\beta$ -R molecules. Such assays may also be used to test LT- $\beta$ -R blocking agents or derivatives of such agents (e.g. fusions, chimeras, mutants, and chemically altered forms) -- alone or in combination -- to optimize the ability of that altered agent to block LT- $\beta$ -R activation.

#### **Production of Soluble LT- $\beta$ -R Molecules**

The LT- $\beta$ -R blocking agents in one embodiment of this invention comprise soluble LT- $\beta$  receptor molecules. **Figure 1** shows the sequence of the extracellular portion of the human LT- $\beta$ -R, which encodes the ligand binding domain. Using the sequence information in **Figure 1** and recombinant DNA techniques well known in the art, functional fragments encoding the LT- $\beta$ -R ligand binding domain can be cloned into a vector and expressed in an appropriate host to produce a soluble LT- $\beta$ -R molecule. Soluble LT- $\beta$ -R molecules that can compete with native LT- $\beta$  receptors for LT ligand binding according to the assays described herein are selected as LT- $\beta$ -R blocking agents.

- 20 -

A soluble LT- $\beta$  receptor comprising amino acid sequences selected from those shown in **Figure 1** may be attached to one or more heterologous protein domains ("fusion domain") to increase the in vivo stability of the receptor fusion protein, or to modulate its biological activity or localization.

Preferably, stable plasma proteins -- which typically have a half-life greater than 20 hours in the circulation -- are used to construct the receptor fusion proteins. Such plasma proteins include but are not limited to: immunoglobulins, serum albumin, lipoproteins, apolipoproteins and transferrin.

Sequences that can target the soluble LT- $\beta$ -R molecule to a particular cell or tissue type may also be attached to the LT- $\beta$ -R ligand binding domain to create a specifically-localized soluble LT- $\beta$ -R fusion protein.

All or a functional portion of the LT- $\beta$ -R extracellular region (**Figure 1**) comprising the LT- $\beta$ -R ligand binding domain may be fused to an immunoglobulin constant region like the Fc domain of a human IgG1 heavy chain (Browning et al., J. Immunol., 154, pp. 33-46 (1995)). Soluble receptor-IgG fusion proteins are preferable, and are common immunological reagents, and methods for their construction are known in the art (see e.g., United States Patent No. 5,225,538 incorporated herein by reference).

A functional LT- $\beta$ -R ligand binding domain may be fused to an immunoglobulin (Ig) Fc domain derived from an immunoglobulin class or subclass other than IgG1. The Fc domains of antibodies belonging to different Ig classes or subclasses can activate diverse secondary effector functions. Activation occurs when the Fc domain is bound by a cognate Fc receptor. Secondary effector functions include the ability to activate the complement system, to cross the placenta, and to bind various microbial proteins. The properties of the different classes and subclasses of immunoglobulins are described in Roitt et al., Immunology, p. 4.8 (Mosby-Year Book Europe Ltd., 3d ed. 1993).

Activation of the complement system initiates cascades of enzymatic reactions that mediate inflammation. The products of the complement system have a variety of functions, including

- 21 -

binding of bacteria, endocytosis, phagocytosis, cytotoxicity, free radical production and solubilization of immune complexes.

The complement enzyme cascade can be activated by the Fc domains of antigen-bound IgG1, IgG3 and IgM antibodies. The Fc domain of IgG2 appears to be less effective, and the Fc domains of IgG4, IgA, IgD and IgE are ineffective at activating complement. Thus one can select a Fc domain based on whether its associated secondary effector functions are desirable for the particular immune response or disease being treated with the LT- $\beta$ -R-Fc fusion protein.

If it would be advantageous to harm or kill the LT ligand-bearing target cell, one could select an especially active Fc domain (IgG1) to make the LT- $\beta$ -R-Fc fusion protein. Alternatively, if it would be desirable to target the LT- $\beta$ -R-Fc fusion to a cell without triggering the complement system, an inactive IgG4 Fc domain could be selected.

Mutations in Fc domains that reduce or eliminate binding to Fc receptors and complement activation have been described (S. Morrison, Annu. Rev. Immunol., 10, pp. 239-65 (1992)). These or other mutations can be used, alone or in combination, to optimize the activity of the Fc domain used to construct the LT- $\beta$ -R-Fc fusion protein.

The production of a soluble human LT- $\beta$ -R fusion protein comprising ligand binding sequences fused to a human immunoglobulin Fc domain (hLT- $\beta$ -R-Fc) is described in Example 1. One CHO line made according to Example 1 that secretes hLT- $\beta$ -R-Fc is called "hLT $\beta$ ;R-hG1 CHO#14". A sample of this line was deposited on July 21, 1995 with the American Type Culture Collection (ATCC) (Rockville, MD) according to the provisions of the Budapest Treaty and was assigned the ATCC accession number CRL11965.

The production of a soluble murine LT- $\beta$ -R fusion molecule (mLT- $\beta$ -R-Fc) is described in Example 2. A CHO line made according to Example 2 that secretes mLT- $\beta$ -R-Fc is called "mLT $\beta$ ;R-hG1 CHO#1.3.BB". A sample of this line was deposited on July 21, 1995 with the American Type Culture Collection (ATCC) (Rockville, MD) according to the provisions of the Budapest Treaty and was assigned the ATCC accession number CRL11964.

- 22 -

All restrictions on the availability to the public of the above ATCC deposits will be irrevocably removed upon the granting of a patent on this application.

Different amino acid residues forming the junction point of the receptor-Ig fusion protein may alter the structure, stability and ultimate biological activity of the soluble LT- $\beta$  receptor fusion protein. One or more amino acids may be added to the C-terminus of the selected LT- $\beta$ -R fragment to modify the junction point with the selected fusion domain.

The N-terminus of the LT- $\beta$ -R fusion protein may also be varied by changing the position at which the selected LT- $\beta$ -R DNA fragment is cleaved at its 5' end for insertion into the recombinant expression vector. The stability and activity of each LT- $\beta$ -R fusion protein may be tested and optimized using routine experimentation and the assays for selecting LT- $\beta$ -R blocking agents described herein.

Using the LT- $\beta$ -R ligand binding domain sequences within the extracellular domain shown in **Figure 1**, amino acid sequence variants may also be constructed to modify the affinity of the soluble LT- $\beta$  receptor or fusion protein for LT ligand. The soluble LT- $\beta$ -R molecules of this invention can compete for surface LT ligand binding with endogenous cell surface LT- $\beta$  receptors. It is envisioned that any soluble molecule comprising a LT- $\beta$ -R ligand binding domain that can compete with cell surface LT- $\beta$  receptors for LT ligand binding is a LT- $\beta$ -R blocking agent that falls within the scope of the present invention.

#### **Soluble LT- $\beta$ -R Molecules As LT- $\beta$ -R Blocking Agents**

A soluble human LT- $\beta$  receptor-immunoglobulin fusion protein (hLT- $\beta$ -R-Fc) was made according to the procedures in Example 1 and tested for its ability to block LT- $\beta$ -R-induced cytotoxicity in human HT29 tumor cells. **Table 1** (Example 3) compares the ability of soluble LT- $\beta$  receptor (hLT- $\beta$ -R-Fc) and TNF receptor (p55-TNF-R-Fc) fusion proteins to block the inhibitory effects of various TNF and soluble LT ligands on HT29 tumor cell growth.



- 23 -

The data in Table 1 indicate the concentrations at which a soluble LT- $\beta$  receptor (hLT- $\beta$ -R-Fc) can block the tumor cell death caused by interaction between LT- $\alpha$ 1/ $\beta$ 2 ligand and cell surface LT- $\beta$  receptors by 50%. The ability to block tumor cell growth at least 20% identifies this soluble LT- $\beta$  receptor as a LT- $\beta$ -R blocking agent according to this invention. As expected, the soluble TNF-R fusion protein (p55-TNF-R-Fc) completely blocked TNF-induced growth inhibition by binding to TNF and preventing its interaction with surface receptor.

The soluble TNF-R fusion protein had no effect on LT ligand (LT- $\alpha$ 1/ $\beta$ 2)-mediated anti-proliferative effects. In contrast, the LT- $\beta$ -R fusion protein blocked LT ligand effects but not the effects of TNF or LT- $\alpha$ . Thus soluble human LT- $\beta$ -R fusion proteins do not interfere with TNF-R activation by TNF and LT- $\alpha$  ligands.

To determine whether LT- $\beta$ -R signalling is also cytotoxic to tumor cells in mice, and whether soluble LT- $\beta$ -R fusion proteins can block LT- $\beta$ -R-induced cytotoxicity, a similar experiment was performed using mouse tumor cells. A soluble murine LT- $\beta$ -R-Fc fusion protein (mLT- $\beta$ -R-Fc; see Example 2) was tested for its ability to block the death of mouse WEHI 164 cells treated with LT ligand (Example 4).

**Figure 2** shows the effects of the soluble murine LT- $\beta$ -R (mLT- $\beta$ -R-Fc) on LT ligand-induced LT- $\beta$ -R signalling in mouse WEHI 164 cells. As this assay indicates, WEHI 164 cells are killed by treatment with soluble LT- $\alpha$ 1/ $\beta$ 2 ligand. Addition of mLT- $\beta$ -R-Fc blocks LT ligand-activated cell death. The control TNF receptor fusion protein (p55TNF-R-Fc) has little effect on blocking cell death.

These data show that a soluble LT- $\beta$ -R fusion protein can effectively compete with surface LT- $\beta$ -R molecules for LT ligand binding. The soluble mLT- $\beta$ -R-Fc fusion protein thus acts as a LT- $\beta$ -R blocking agent in mice.

- 24 -

**Source of Anti-Human LT- $\beta$ -R Antibodies**

In another embodiment of this invention, antibodies directed against the human LT- $\beta$  receptor (anti-LT- $\beta$ -R Abs) function as LT- $\beta$ -R blocking agents. The anti-LT- $\beta$ -R Abs of this invention can be polyclonal or monoclonal (mAbs) and can be modified to optimize their ability to block LT- $\beta$ -R signalling, their in vivo bioavailability, stability, or other desired traits.

Polyclonal antibody sera directed against the human LT- $\beta$  receptor are prepared using conventional techniques by injecting animals such as goats, rabbits, rats, hamsters or mice subcutaneously with a human LT- $\beta$  receptor-Fc fusion protein (Example 1) in complete Freund's adjuvant, followed by booster intraperitoneal or subcutaneous injection in incomplete Freund's. Polyclonal antisera containing the desired antibodies directed against the LT- $\beta$  receptor are screened by conventional immunological procedures.

Mouse monoclonal antibodies (mAbs) directed against a human LT- $\beta$  receptor-Fc fusion protein are prepared as described in Example 5. A hybridoma cell line (BD.A8.AB9) which produces the mouse anti-human LT- $\beta$ -R mAb BDA8 was deposited on January 12, 1995 with the American Type Culture Collection (ATCC) (Rockville, MD) according to the provisions of the Budapest Treaty, and was assigned the ATCC accession number HB11798. All restrictions on the availability to the public of the above ATCC deposits will be irrevocably removed upon the granting of a patent on this application.

Various forms of anti-LT- $\beta$ -R antibodies can also be made using standard recombinant DNA techniques (Winter and Milstein, Nature, 349, pp. 293-99 (1991)). For example, "chimeric" antibodies can be constructed in which the antigen binding domain from an animal antibody is linked to a human constant domain (e.g. Cabilly et al., US 4,816,567; Morrison et al., Proc. Natl. Acad. Sci. U.S.A., 81, pp. 6851-55 (1984)). Chimeric antibodies reduce the observed immunogenic responses elicited by animal antibodies when used in human clinical treatments.

- 25 -

In addition, recombinant "humanized antibodies" which recognize the LT- $\beta$ -R can be synthesized. Humanized antibodies are chimeras comprising mostly human IgG sequences into which the regions responsible for specific antigen-binding have been  
5 inserted (e.g. WO 94/04679). Animals are immunized with the desired antigen, the corresponding antibodies are isolated, and the portion of the variable region sequences responsible for specific antigen binding are removed. The animal-derived antigen binding regions are then cloned into the appropriate  
10 position of human antibody genes in which the antigen binding regions have been deleted. Humanized antibodies minimize the use of heterologous (inter-species) sequences in human antibodies, and are less likely to elicit immune responses in the treated subject.

15 Construction of different classes of recombinant anti-LT- $\beta$ -R antibodies can also be accomplished by making chimeric or humanized antibodies comprising the anti-LT- $\beta$ -R variable domains and human constant domains (CH1, CH2, CH3) isolated from different classes of immunoglobulins. For example, anti-LT- $\beta$ -R  
20 IgM antibodies with increased antigen binding site valencies can be recombinantly produced by cloning the antigen binding site into vectors carrying the human  $\mu$  chain constant regions (Arulanandam et al., J. Exp. Med., 177, pp. 1439-50 (1993); Lane et al., Eur. J. Immunol., 22, pp. 2573-78 (1993); Traunecker et  
25 al., Nature, 339, pp. 68-70 (1989)).

In addition, standard recombinant DNA techniques can be used to alter the binding affinities of recombinant antibodies with their antigens by altering amino acid residues in the vicinity of the antigen binding sites. The antigen  
30 binding affinity of a humanized antibody can be increased by mutagenesis based on molecular modelling (Queen et al., Proc. Natl. Acad. Sci. U.S.A., 86, pp. 10029-33 (1989); WO 94/04679).

It may be desirable to increase or to decrease the affinity of anti-LT- $\beta$ -R Abs for the LT- $\beta$ -R depending on the  
35 targeted tissue type or the particular treatment schedule envisioned. For example, it may be advantageous to treat a patient with constant levels of anti-LT- $\beta$ -R Abs with reduced ability to signal through the LT- $\beta$  pathway for semi-prophylactic

- 26 -

treatments. Likewise, inhibitory anti-LT- $\beta$ -R Abs with increased affinity for the LT- $\beta$ -R may be advantageous for short-term treatments.

#### **Anti-LT- $\beta$ -R Antibodies As LT- $\beta$ -R Blocking Agents**

5           Anti-LT- $\beta$ -R antibodies that act as LT- $\beta$ -R blocking agents may be selected by testing their ability to inhibit LT- $\beta$ -R-induced cytotoxicity in tumor cells (Example 5).

          In a preferred embodiment of this invention, compositions and methods comprise the mouse anti-human LT- $\beta$ -R  
10 mAb BDA8. **Figure 3** shows that mAb BDA8 acts as a LT- $\beta$ -R blocking agent as defined by this invention. WiDr tumor cells stop growing in the presence of IFN- $\gamma$  and soluble LT- $\alpha$ 1/ $\beta$ 2 ligand. Control antibodies (IgG1) have no effect on this growth inhibition. In contrast, the anti-LT- $\beta$ -R mAb BDA8 blocks the  
15 ability of soluble LT- $\alpha$ 1/ $\beta$ 2 ligand to inhibit WiDr cell growth. Thus an antibody directed against human LT- $\beta$ -R can function as a LT- $\beta$ -R blocking agent as defined by the present invention.

          By testing other antibodies directed against the human LT- $\beta$  receptor, it is expected that additional anti-LT- $\beta$ -R  
20 antibodies that function as LT- $\beta$ -R blocking agents in humans can be identified using routine experimentation and the assays described herein.

#### **Source of Anti-Surface LT Ligand Antibodies**

          Another preferred embodiment of this invention  
25 involves compositions and methods which comprise antibodies directed against LT ligand that function as LT- $\beta$ -R blocking agents. As described above for the anti-LT- $\beta$ -R Abs, anti-LT ligand antibodies that function as LT- $\beta$ -R blocking agents can be polyclonal or monoclonal, and can be modified according to  
30 routine procedures to modulate their antigen binding properties and their immunogenicity.

          The anti-LT antibodies of this invention can be raised against either one of the two LT subunits individually, including soluble, mutant, altered and chimeric forms of the LT  
35 subunit. If LT subunits are used as the antigen, preferably they are LT- $\beta$  subunits. If LT- $\alpha$  subunits are used, it is

- 27 -

preferred that the resulting anti-LT- $\alpha$  antibodies bind to surface LT ligand and do not cross-react with secreted LT- $\alpha$  or modulate TNF-R activity (according to the assays described in Example 3).

5 Alternatively, antibodies directed against a homomeric (LT- $\beta$ ) or a heteromeric (LT- $\alpha/\beta$ ) complex comprising one or more LT subunits can be raised and screened for activity as LT- $\beta$ -R blocking agents. Preferably, LT- $\alpha 1/\beta 2$  complexes are used as the antigen. As discussed above, it is preferred that the resulting  
10 anti-LT- $\alpha 1/\beta 2$  antibodies bind to surface LT ligand without binding to secreted LT- $\alpha$  and without affecting TNF-R activity.

The production of polyclonal anti-human LT- $\alpha$  antibodies is described in applicants' co-pending application (WO 94/13808). Monoclonal anti-LT- $\alpha$  and anti-LT- $\beta$  antibodies  
15 have also been described (Browning et al., J. Immunol., 154, pp. 33-46 (1995)).

Mouse anti-human LT- $\beta$  mAbs were prepared as described in Example 6. A hybridoma cell line (B9.C9.1) which produces the mouse anti-human LT- $\beta$ -R mAb B9 was deposited on July 21,  
20 1995 with the American Type Culture Collection (ATCC) (Rockville, MD) according to the provisions of the Budapest Treaty, and was assigned the ATCC accession number HB11962.

Monoclonal hamster anti-mouse LT- $\alpha/\beta$  antibodies were prepared as described in Example 7. A hybridoma cell line  
25 (BB.F6.1) which produces the hamster anti-mouse LT- $\alpha/\beta$  mAb BB.F6 was deposited on July 21, 1995 with the American Type Culture Collection (ATCC) (Rockville, MD) according to the provisions of the Budapest Treaty, and was assigned the ATCC accession number HB11963.

30 All restrictions on the availability to the public of the above ATCC deposits will be irrevocably removed upon the granting of a patent on this application.

#### **Anti-LT Ligand Antibodies As LT- $\beta$ -R Blocking Agents**

A fluorescence-activated cell sorting (FACS) assay was  
35 developed to screen for antibodies directed against LT subunits and LT complexes that can act as LT- $\beta$ -R blocking agents (Examples 6 and 7). In this assay, soluble human LT- $\beta$ -R-Fc

- 28 -

fusion protein is added to PMA-activated II-23 cells -- which express surface LT complexes (Browning et al., J. Immunol., 154, pp. 33-46 (1995)) -- in the presence of increasing amounts of the test antibody. An antibody that can inhibit LT- $\beta$  receptor-ligand interaction by at least 20% is selected as a LT- $\beta$ -R blocking agent.

The results of this assay performed to test the mouse anti-human LT- $\beta$  mAb B9 are shown in **Figure 4**. **Figure 4** shows that anti-LT- $\beta$  mAb B9 can selectively block the binding of soluble LT- $\beta$ -R-Fc fusion proteins to surface LT ligands induced on activated cells. These results confirm that antibodies directed against a LT ligand subunit will function as an LT- $\beta$ -R blocking agent.

The FACS assay described above was also used to test mAbs raised in hamster against a soluble mouse LT- $\alpha/\beta$  complex (Example 7). The results of this assay performed to test the hamster anti-mouse LT- $\alpha/\beta$  mAb BB.F6 are shown in Table 2 (Example 7). Table 2 shows that anti-LT- $\alpha/\beta$  mAb BB.F6 can effectively block the binding of soluble mLT- $\beta$ -R-Fc fusion proteins (Example 2) to surface LT ligands expressed on a murine T cell hybridoma and is thus a LT- $\beta$ -R blocking agent according to this invention.

Using a LT- $\alpha/\beta$  complex rather than a LT subunit as an antigen to immunize an animal may lead to more efficient immunization, or may result in antibodies having higher affinities for surface LT ligand. It is conceivable that by immunizing with the LT- $\alpha/\beta$  complex, antibodies which recognize amino acid residues on both the LT- $\alpha$  and the LT- $\beta$  subunits (e.g., residues that form an LT- $\alpha/\beta$  cleft) can be isolated. By testing antibodies directed against human LT- $\alpha/\beta$  heteromeric complexes, it is expected that additional anti-LT antibodies that function as LT- $\beta$ -R blocking agents in humans can be identified using routine experimentation and the assays described herein.

### LT- $\beta$ -R Blocking Agents Inhibit Th1 Cell-Mediated Contact Hypersensitivity in Mouse

- 29 -

The LT- $\beta$ -R blocking agents of this invention can inhibit Th1 cell-mediated immune responses. One such Th1-mediated response is delayed type hypersensitivity (DTH; Cher and Mosmann, J. Immunol., 138, pp. 3688-94 (1987); see also I. Roitt et al., Immunology, pp. 22.1-22.12, Mosby-Year Book Europe Ltd., 3d ed. (1993) for a general discussion). DTH is evoked when antigen-sensitized Th1 cells secrete cytokines following a secondary contact with the same antigen. The Th1 cytokines attract and activate macrophages that release additional effector molecules which trigger inflammatory reactions.

DTH reactions are classified into three different types: contact hypersensitivity, tuberculin-type hypersensitivity and granulomatous reactions. The three types of hypersensitivity (HS) may be distinguished by the speed and nature of the response to foreign antigen when it is applied directly to or injected beneath the skin of a sensitized subject. The DTH reaction is monitored by measuring the rate and degree to which the skin thickens.

Tuberculin-type HS reactions are skin reactions which occur at the injection site of a foreign antigen from a microorganism to which the subject has been previously exposed (e.g. mycobacterium tuberculosis or M. leprae). This skin reaction, which is maximal between 48 and 72 hours, is frequently used as the basis for diagnostic sensitivity tests to previously-encountered microorganisms (e.g. the tuberculin skin test). As a tuberculin-type lesion develops, it can become a granulomatous reaction if the antigen persists in the tissue.

Granulomatous reactions are clinically the most serious DTH reactions because they can lead to many of the pathological effects associated with Th1 cell-mediated diseases. Granulomatous reactions occur when antigens or immune complexes fail to clear from macrophages and continue to stimulate Th1 cytokine secretion. Chronic inflammation and aggregation of activated macrophages at the site of the stimulus characterize granulomatous reactions.

A core of epithelial cells and macrophages, which can also be surrounded by lymphocytes and fibrotic depositions, form a hardened structure called a granuloma. Sometimes there is

- 30 -

extensive cell death in the core of the granuloma (e.g. in tuberculosis-affected lung tissue). Hardening in the target tissue of a granulomatous reaction occurs in about 4 weeks.

Agents which affect the frequency of granuloma formation can be identified using schistosome-infected mice (Amiri et al., Nature, 356, pp. 604-607 (1992)). Schistosome worms (blood flukes) can cause a parasitic disease leading to granuloma formation around the schistosome eggs deposited in portal venules of the infected liver. Agents that inhibit this Th1 cell-mediated DTH response may decrease the size of the granuloma, or the frequency or rate of granuloma formation in schistosome-infected mouse livers. Cellular reaction to the schistosome eggs can be assessed by quantitating the number and size of granulomas formed in mice treated with increasing concentrations of a putative LT- $\beta$ -R blocking agent over time.

Contact hypersensitivity (CHS) is a class of DTH in which skin is the target organ. In CHS, an inflammatory response is caused by locally applying a reactive hapten onto the skin. Allergens generally comprise at least one hapten molecule, which is usually too small to be antigenic on its own. The hapten penetrates the epidermis and reacts with a normal protein under the skin to produce a novel antigenic complex.

Re-exposure of a sensitized subject to the hapten triggers the DTH response. The hapten-carrier protein conjugate, in combination with antigen presenting cells, activates effector mechanisms that trigger the release of cytokines (including IL-2, IL-3, IFN- $\gamma$  and GM-CSF). The cascade of released cytokines causes CD4+ T cells to proliferate, the expression patterns of various cell surface adhesion molecules to change, and the attraction of T cells and macrophages to the skin at the site of inflammation. The cytokine cascade and resulting vasodilation, cellular infiltration and edema of the dermis and epidermis leads to swelling and inflammation of the target tissue, which accounts for the measurable skin thickening in response to DTH reactions.

The degree to which a particular hapten can sensitize an individual depends on a variety of factors. These factors



- 31 -

include how well the hapten can penetrate the skin and react with a host carrier protein to form a conjugate. One hapten that sensitizes nearly all individuals is 2,4-dinitro-fluorobenzene (DNFB).

- 5           The skin CHS response to a hapten such as DNFB is a classic animal model for cell-mediated immunity. Localization of this CHS response to the ear of a sensitized mouse allows easy, accurate and reproducible quantitation of this cell-mediated immune response in vivo by measuring ear thickness.
- 10 The details of the murine CHS reaction and the histopathology of the DNFB-induced inflammatory response have been reported (Chisholm et al., Eur. J. Immunol., 23, pp. 682-688 (1993)).

- The ability of DNFB to induce a contact hypersensitivity response in most individuals can be used to
- 15 identify agents that reduce or eliminate the inflammatory responses associated with Th1 cell-mediated DTH reactions. A soluble murine LT- $\beta$ -R-Fc fusion protein effectively inhibits DNFB-induced contact hypersensitivity responses in mice (Example 8). Mice were initially sensitized by applying DNFB onto the
- 20 bottom of each hind foot on two consecutive days. Five days after the initial sensitization, a sub-irritant dose of DNFB in carrier solution was applied to the surfaces of the left ear. Carrier solution alone was applied to the right ear as a control.

- 25           Increasing concentrations of the LT- $\beta$ -R blocking agent mLT- $\beta$ -R-Fc (Example 2) were then injected intravenously into the mice (Example 8). Injections of PBS buffer alone, or of a human IgG fusion protein (LFA3-Fc) served as negative controls, and injection of an anti-VLA4-specific mAb (PS/2 mAb) known to
- 30 inhibit CHS served as a positive control. Twenty-four hours after challenge, the thickness of each ear (DNFB-challenged and -unchallenged) was measured. Inhibition of the ear swelling response by the LT- $\beta$ -R blocking agent was judged by comparison of treated groups with their negative control group.

- 35           **Figure 5** shows that mLT- $\beta$ -R-Fc causes a significant reduction in the ear swelling response of DNFB-treated mice compared to uninhibited DNFB-treated control animals (PBS and LFA3-Fc). Soluble LT- $\beta$ -R can block this CHS reaction as

- 32 -

effectively as the inhibitor anti-VLA4-specific mAb (PS/2 mAb), which acts by blocking the influx of T cells into the challenge site (Chisholm et al., Eur. J. Immunol., 23, pp. 682-88 (1993)).

These data show that a soluble LT- $\beta$ -R fusion protein which acts as a LT- $\beta$ -R blocking agent in vitro can also effectively inhibit a Th1 cell-mediated immune response when administered to an animal. The LT- $\beta$ -R blocking agents of this invention identified in vitro can be tested using this ear swelling assay, or other DTH assays such as those described above, to select additional LT- $\beta$ -R blocking agents that will be useful for reducing the severity of Th1 cell-associated immune responses in vivo.

**LT- $\beta$ -R Blocking Agents Do Not Inhibit a Th2 Cell-Mediated (Humoral) Immunological Response**

As shown above, the LT- $\beta$ -R blocking agents of this invention can inhibit a Th1 cell-mediated effector mechanism such as contact delayed type hypersensitivity (**Figure 5**). This Th1 cell-mediated response is inhibited without significantly affecting Th2 cell-dependent responses. The differential effect of LT- $\beta$ -R blocking agents on Th1 cell-mediated immune responses was shown by monitoring a Th2-cell dependent immune response -- such as a primary antibody response and isotype switching -- in the presence of an LT- $\beta$ -R blocking agent.

Mice were injected five times over the course of a ten day period with either soluble LT- $\beta$ -R fusion protein (mLT- $\beta$ -R-Fc; Example 2) or control IgG fusion protein (LFA3-Fc), or were left untreated. After the second injection, all mice were injected in the base of the tail with 100  $\mu$ l of complete Freund's adjuvant containing 100  $\mu$ g of ovalbumin. After 11 days, primary serum anti-ovalbumin-specific antibody titers were analyzed using an ELISA specific for IgG1, IgG2a and IgM isotypes.

**Figure 6** shows the effect of the mouse LT- $\beta$ -R blocking agent mLT- $\beta$ -R-Fc on serum anti-ovalbumin antibody production in mice immunized with ovalbumin (Example 9). Administering the

- 33 -

LT- $\beta$ -R blocking agent does not significantly affect primary antibody titers following ovalbumin immunization. By comparison, interfering with CD40 ligand-induced CD40 receptor signalling completely blocks the antigen-specific IgG response in mice (Renshaw et al., J. Exp. Med., 180, pp. 1889-1900 (1994)). CD40 is another ligand/receptor pair in the TNF family.

Total immunoglobulin production and maturation is clearly Th2 cell-dependent. However, there is also evidence that the Th1 cytokine IFN- $\gamma$  participates but is not absolutely required for the switch to the IgG2a subclass (Huang et al., Science, 259, pp. 1742-45 (1993)). The LT- $\beta$ -R blocking agent mLT- $\beta$ -R-Fc did not inhibit the IgG2a switch in these experiments. It is possible that the LT- $\beta$ -R blocking agents of this invention do not block this humoral aspect of a Th1 cell-mediated response. In addition, the proliferatory responses of lymphocytes from the mLT- $\beta$ -R-Fc-treated mice were not decreased (Example 10; **Figure 7**).

These experiments indicate that a therapy based on administering the LT- $\beta$ -R blocking agents of this invention will not adversely affect Th2 dependent antibody production functions of an immune response. The normal pattern of antibody response illustrated in **Figure 6** also indicates that an intensive treatment with soluble mLT- $\beta$ -R-Fc was not toxic to the mice, further indicating the useful therapeutic nature of the compositions and methods set forth in this invention.

#### **T Helper Cell-Mediated Diseases**

Many organ-specific autoimmune conditions appear to involve pathological Th1 response. These data have been reviewed (Modlin and Nutman, Current Opinion in Immunol., 5, pp. 511-17 (1993); Romagnani et al., Ann. Rev. Immunol., 12, pp. 227-57 (1994)). These organ-specific autoimmune conditions include: multiple sclerosis, insulin-dependent diabetes, sympathetic ophthalmia, uveitis and psoriasis.

- 34 -

Insulin-dependent diabetes mellitus is an autoimmune disease in which the insulin-producing beta pancreatic cells are destroyed by leukocytes infiltrating into the islets of Langerhans. Diabetes can be rapidly induced in neonatal nonobese diabetic (NOD) mice by transferring activated prediabetic splenocytes. Recently, Th1- or Th2-like cells, otherwise genetically similar, were transferred into neonatal NOD mice. Only the Th1 cells rapidly induced diabetes -- and in almost all recipients (Katz et al., Science, 268, pp. 1185-88 (1995)). This indicates that the LT- $\beta$ -R blocking agents of this invention -- which can inhibit the effects of a Th1 cell-mediated immune response in vivo -- will be useful for treating or preventing insulin-dependent diabetes.

Several systemic autoimmune diseases, including various arthritides, are Th1 cell-associated. Rheumatoid arthritis and Sjorgren's syndrome both appear to involve Th0 and Th1 cells. In contrast, systemic lupus erythematosus (SLE) appears to have an aberrant Th0/Th2 dominated response.

Some chronic inflammatory diseases also appear to have an aberrant Th1 type response, including inflammatory bowel disease, sarcoidosis of the lung and allograft rejection. Inflammatory bowel disease (IBD) in humans encompasses at least two categories, ulcerative colitis and Crohn's disease. Both disorders are believed to result from immunopathologic autoimmune like disorders. In some mouse models of IBD, it is clear that some agents that block Th1 responses can block the development or course of the disease (F. Powrie et al, Immunity 1:553 1994). It is possible that inhibition of the Th1 component of the immune response would have beneficial effects in human IBD. Many models of IBD have been described and have been reviewed (C. Elson et al, Gastroenterology 109:1344 1995). There are at least three groups of models, chemically induced, polymer/microbial-induced and immunological types using mutant mice.

In one commonly used polymer/microbial-induced model, dextran sulphate solution is introduced into the drinking water of mice and upon ingestion, the epithelial lining of the gut is irritated leading to a profound immune response to the damage.

- 35 -

The animals develop colitis which is manifested as diarrhea, blood in the stool, loss of body weight and a shortening of the colon length due to expansion of the colon wall. This model induces a left-sided colitis and epithelial dysplasia which can lead to cancer which are features of ulcerative colitis.

A second model consists of transplanting a selected set of CD4 T cells into a scid mouse, i.e. a mouse lacking T and B cells (F. Powrie et al International Immunology 5:1461-1471 1993; Morrissey et al, J. Exp. Med. 178:237 1993). As the selected cells, called CD45RB<sup>hi</sup> cells expand and reconstitute the scid mouse, the normal mechanisms preventing the appearance of autoreactive T cells are dysfunctional and autoreactive cells develop. In rats, cells reactive with many organs are observed whereas, in the mouse, the reactivity occurs primarily in the bowel. Agents which either alter the way the autoreactive cells expand and develop or agents which can block the ability of the cells to attack the bowel will have efficacy in this model. Moreover, as this model at least partially mimics the pathological development of autoreactive immune system cells, treatments that block this model may actually have disease modifying behavior in humans. In this model, antibodies to TNF can block disease (F. Powrie et al Immunity 1, 552 1994) and these antibodies have been found to be efficacious in the treatment of human disease (H. M. van Dulleman et al. Gastroenterology 109:109 1995). Thus, this model can forecast which agents may be therapeutically useful in IBD. Moreover, as the CD45RB model is an example of a Th1 mediated disease process and indeed in rats, the model leads to disease in many organs, the efficacy of LT $\beta$ R-Ig in this system indicates that LT $\beta$ R-Ig or other means of blocking the LT $\beta$ R interactions with its ligand may be beneficial in a wide range of related immunological diseases.

In general, the exact contribution of auto-antibodies versus specific T cells has not been delineated in these autoimmune diseases. Cellular responses may make major contributions to pathogenicity in those systemic autoimmune diseases currently thought to be primarily antibody driven, e.g. the various arthritides.

- 36 -

The normal immune response to some pathogenic infectious agents also elicits a Th1 response that can become excessive and present itself as a medical problem. Examples of granulomatous reactions (a class of DTH response described above) that lead to severe medical problems include leprosy, granuloma formation in the lungs of tuberculosis patients, sarcoidosis and schistosomiasis (Roitt et al., Immunology, pp. 22.5-6 (Mosby-Year Book Europe Ltd., 3d ed. 1993)). Psoriasis is also likely to be mediated by Th1 cells.

Cytolytic T cells, i.e. CTLs (CD8 positive T cells) may also subdivide into Th1- and Th2-like populations. Therefore it is possible that much of what is known regarding the Th groups will also apply to CD8+ cells, which are primarily involved in anti-viral and grafted tissue rejection responses.

#### 15 **Treatments Using LT- $\beta$ -R Blocking Agents**

The compositions of this invention will be administered at an effective dose to treat the particular clinical condition addressed. Determination of a preferred pharmaceutical formulation and a therapeutically efficient dose regiment for a given application is well within the skill of the art taking into consideration, for example, the condition and weight of the patient, the extent of desired treatment and the tolerance of the patient for the treatment. Doses of about 1 mg/kg of a soluble LT- $\beta$ -R are expected to be suitable starting points for optimizing treatment doses.

Determination of a therapeutically effective dose can also be assessed by performing in vitro experiments that measure the concentration of the LT- $\beta$ -R blocking agent required to coat target cells (LT- $\beta$ -R or LT ligand-positive cells depending on the blocking agent) for 1 to 14 days. The receptor-ligand binding assays described herein can be used to monitor the cell coating reaction. LT- $\beta$ -R or LT ligand-positive cells can be separated from activated lymphocyte populations using FACS. Based on the results of these in vitro binding assays, a range of suitable LT- $\beta$ -R blocking agent concentrations can be selected to test in animals according to the assays described herein.

- 37 -

Administration of the soluble LT- $\beta$ -R molecules, anti-LT ligand and anti-LT- $\beta$ -R Abs of this invention, alone or in combination, including isolated and purified forms of the antibodies or complexes, their salts or pharmaceutically acceptable derivatives thereof, may be accomplished using any of the conventionally accepted modes of administration of agents which exhibit immunosuppressive activity.

The pharmaceutical compositions used in these therapies may also be in a variety of forms. These include, for example, solid, semi-solid and liquid dosage forms such as tablets, pills, powders, liquid solutions or suspensions, suppositories, and injectable and infusible solutions. The preferred form depends on the intended mode of administration and therapeutic application. Modes of administration may include oral, parenteral, subcutaneous, intravenous, intralesional or topical administration.

The soluble LT- $\beta$ -R molecules, anti-LT ligand and anti-LT- $\beta$ -R Abs of this invention may, for example, be placed into sterile, isotonic formulations with or without cofactors which stimulate uptake or stability. The formulation is preferably liquid, or may be lyophilized powder. For example, the soluble LT- $\beta$ -R molecules, anti-LT ligand and anti-LT- $\beta$ -R Abs of this invention may be diluted with a formulation buffer comprising 5.0 mg/ml citric acid monohydrate, 2.7 mg/ml trisodium citrate, 41 mg/ml mannitol, 1 mg/ml glycine and 1 mg/ml polysorbate 20. This solution can be lyophilized, stored under refrigeration and reconstituted prior to administration with sterile Water-For-Injection (USP).

The compositions also will preferably include conventional pharmaceutically acceptable carriers well known in the art (see for example Remington's Pharmaceutical Sciences, 16th Edition, 1980, Mac Publishing Company). Such pharmaceutically acceptable carriers may include other medicinal agents, carriers, genetic carriers, adjuvants, excipients, etc., such as human serum albumin or plasma preparations. The compositions are preferably in the form of a unit dose and will usually be administered one or more times a day.

- 38 -

The pharmaceutical compositions of this invention may also be administered using microspheres, liposomes, other microparticulate delivery systems or sustained release formulations placed in, near, or otherwise in communication with affected tissues or the bloodstream. Suitable examples of sustained release carriers include semipermeable polymer matrices in the form of shaped articles such as suppositories or microcapsules. Implantable or microcapsular sustained release matrices include polylactides (U.S. Patent No. 3,773,319; EP 58,481), copolymers of L-glutamic acid and ethyl-L-glutamate (Sidman et al., Biopolymers, 22, pp. 547-56 (1985)); poly(2-hydroxyethyl-methacrylate) or ethylene vinyl acetate (Langer et al., J. Biomed. Mater. Res., 15, pp. 167-277 (1981); Langer, Chem. Tech., 12, pp. 98-105 (1982)).

Liposomes containing soluble LT- $\beta$ -R molecules, anti-LT ligand and anti-LT- $\beta$ -R Abs of this invention, alone or in combination, can be prepared by well-known methods (See, e.g. DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. U.S.A., 82, pp. 3688-92 (1985); Hwang et al., Proc. Natl. Acad. Sci. U.S.A., 77, pp. 4030-34 (1980); U.S. Patent Nos. 4,485,045 and 4,544,545). Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol.% cholesterol. The proportion of cholesterol is selected to control the optimal rate of soluble LT- $\beta$ -R molecule, anti-LT ligand and anti-LT- $\beta$ -R Ab release.

The soluble LT- $\beta$ -R molecules, anti-LT ligand and anti-LT- $\beta$ -R Abs of this invention may also be attached to liposomes containing other LT- $\beta$ -R blocking agents, immunosuppressive agents or cytokines to modulate the LT- $\beta$ -R blocking activity. Attachment of LT- $\beta$ -R molecules, anti-LT ligand and anti-LT- $\beta$ -R Abs to liposomes may be accomplished by any known cross-linking agent such as heterobifunctional cross-linking agents that have been widely used to couple toxins or chemotherapeutic agents to antibodies for targeted delivery. Conjugation to liposomes can also be accomplished using the carbohydrate-directed cross-linking reagent 4-(4-maleimidophenyl) butyric acid hydrazide (MPBH) (Duzgunes et al., J. Cell. Biochem. Abst. Suppl. 16E 77 (1992)).



- 39 -

**Advantages of therapeutic compositions comprising  
LT- $\beta$ -R blocking agents**

The LT- $\beta$ -R blocking agents of this invention are capable of selectively inhibiting Th1 and not Th2 cell-dependent immune effector mechanisms. LT- $\beta$ -R blocking agents will be useful in treating conditions that are exacerbated by the activities of Th1-type cytokines (e.g., IL-2 and IFN- $\gamma$ ). Because Th1 cytokines can inhibit Th2 cell-dependent responses, LT- $\beta$ -R blocking agents may also indirectly stimulate certain Th2 cell-dependent responses that are normally inhibited by Th1-induced cytokine cascades.

The ability to selectively suppress Th1 (or indirectly stimulate Th2) cell responses will be useful for treating abnormalities in diverse cell-mediated immune responses including various autoimmune and chronic inflammatory conditions, antigen tolerance, and cellular rejection of tissue grafts and organ transplants.

As discussed above, treatment of Th1 cell-based immunological conditions generally employs immunomodulatory and immunosuppressive agents which have pleiotropic effects on a wide variety of cell types and immunological responses. These non-specific immunosuppressive agents are generally required in high and often cytotoxic doses that cause adverse side effects.

The ability to shift the character of an immunological response is supported in the recent study of mouse diabetes discussed above (Katz et al., Science, 268, pp. 1185-88 (1995)), and in an allogeneic transplant model (Sayegh et al., J. Exp. Med., 181, pp. 1869-74 (1995)). In the latter study, a fusion protein that blocks the CD28-B7 T cell costimulatory pathway was shown to induce renal graft tolerance. The tolerance correlated with a decrease in Th1 cytokines and an increase in Th2 cytokines in vivo. These data indicate that the LT- $\beta$ -R blocking agents of this invention will be useful in suppressing cellular rejection of tissue grafts and organ transplants by inhibiting Th1 cell-mediated cytokine release.

The LT- $\beta$ -R blocking agents of the compositions and methods of this invention can be modified to obtain a desirable level of LT- $\beta$ -R signalling depending on the condition, disorder

- 40 -

or disease being treated. It is envisioned that the absolute level of LT- $\beta$ -R signalling can be fine-tuned by manipulating the concentration and the affinities of the LT- $\beta$ -R blocking agents for their respective molecular targets.

5           For example, in one embodiment of this invention, compositions comprising soluble LT- $\beta$ -R molecules are administered to a subject. The soluble LT- $\beta$  receptor can effectively compete with cell surface LT- $\beta$  receptors for binding surface LT ligands. The ability to compete with surface LT  
10 ligands depends on the relative concentrations of the soluble and the cell surface LT- $\beta$ -R molecules, and on their relative affinities for ligand binding.

Soluble LT- $\beta$ -R molecules harboring mutations that increase or decrease the binding affinity of that mutant soluble  
15 LT- $\beta$ -R with surface LT ligand can be made using standard recombinant DNA techniques well known to those of skill in the art. Large numbers of molecules with site-directed or random mutations can be tested for their ability to act as LT- $\beta$ -R blocking agents using routine experimentation and the techniques  
20 described herein.

Similarly, in another embodiment of this invention, antibodies directed against either the LT- $\beta$  receptor or one or more of the LT ligand subunits function as LT- $\beta$ -R blocking agents. The ability for these antibodies to block LT- $\beta$  receptor  
25 signalling can be modified by mutation, chemical modification or by other methods that can vary the effective concentration or activity of the antibody delivered to the subject.

The ability to diminish LT- $\beta$ -R signalling without completely inhibiting it may be important for establishing or  
30 maintaining reduced levels of LT- $\beta$ -R signalling that support normal immune function while inhibiting Th1-cell mediated responses which are exaggerated or abnormal.

Disruption of the LT- $\alpha$  gene in a mouse leads to aberrant peripheral lymphoid organ development (De Togni et al.,  
35 Science, 264, pp. 703-7 (1994)). Such mice lacked lymph nodes and their spleens lacked the usually clear demarcation between T and B cell-rich regions in the follicles. We believe that this phenotype is associated with loss of surface LT-induced LT- $\beta$ -R

- 41 -

signalling because similar phenotypes have not been observed by modulating TNF-R activity. The ability to selectively or to partially block the LT- $\beta$ -R pathway may thus be useful in treating abnormal lymphoid organ development associated with mis- or over-expression of signalling by the LT- $\beta$ -R pathway.

Some Th1-associated reactions are critical components of a number of cell-mediated immune responses (Romagnani, S., Ann. Rev. Immunol., 12, pp. 227-57 (1994)), and absolute inhibition of Th1 cell activity may not be desirable in certain circumstances. For example, a mouse can effectively resist a parasitic infection when a good Th1 response can be mounted. Infectious agents such as Listeria and Toxoplasma also elicit strong Th1-type responses. In humans, mycobacterium tuberculosis responses appear to be Th1-based. Leishmaniasis pathogenicity correlates with responses similar to the Th1 responses characterized in mouse (Reed and Scott, Current Opinion in Immunol., 5, pp. 524-31 (1993)).

The ability to influence the level of Th1 inhibition by blocking LT- $\beta$ -R signalling may be important in maximizing the beneficial results which can be achieved by treatments with the LT- $\beta$ -R blocking agents of this invention.

The following are examples which illustrate the soluble LT- $\beta$  receptors, anti-LT ligand and anti-LT- $\beta$ -R antibodies of this invention and the methods used to characterize them. These examples should not be construed as limiting: the examples are included for purposes of illustration and the present invention is limited only by the claims.

#### EXAMPLE 1

30

##### **Preparation of Soluble Human LT- $\beta$ Receptors as Immunoglobulin Fc Fusion Proteins**

The sequence of a human cDNA clone isolated from a library of human 12p transcribed sequences derived from a somatic cell hybrid (Baens et al., Genomics, 16, pp. 214-18 (1993)), was entered into GenBank and was later identified as the sequence which encodes human LT- $\beta$ -R. The sequence of this

- 42 -

full-length human LT- $\beta$ -R cDNA clone has been available since 1992 as GenBank entry L04270.

The extracellular domain of LT- $\beta$ -R up to the transmembrane region (**Figure 1**) was amplified by PCR from a cDNA clone using primers that incorporated NotI and SalI restriction enzyme sites on the 5' and 3' ends, respectively (Browning et al., J. Immunol., 154, pp. 33-46 (1995)). The amplified product was cut with NotI and SalI, purified and ligated into a NotI-linearized vector pMDR901 along with a SalI-NotI fragment encoding the Fc region of human IgG1. The resultant vector contained the dihydrofolate reductase gene and the LT- $\beta$ -R-Fc fusion protein driven by separate promoters.

The vector was electroporated into CHO dhfr<sup>-</sup> cells and methotrexate-resistant clones were isolated as per standard procedures. The LT- $\beta$ -R-Fc was secreted into the medium and an ELISA assay was used to select for cell lines producing the highest level of the receptor fusion protein. A high-producing cell line was grown to large numbers and the conditioned medium collected. The pure LT- $\beta$  receptor fusion protein was isolated by Protein A Sepharose Fast Flow affinity chromatography (Pharmacia).

## EXAMPLE 2

### Preparation of Soluble Murine LT- $\beta$ Receptors as Immunoglobulin Fc Fusion Proteins

A complete cDNA clone of the mLT- $\beta$ -R was prepared by ligating a 5' NotI/ApaLI and 3' ApaLI/NotI fragments from two partial cDNA isolates into the NotI site of pCDNA3 (InVitrogen, San Diego, CA). The sequence of this cDNA clone is accessible as GenBank entry U29173. No coding sequence differences were noted when compared with another sequence entry for mLT- $\beta$ -R found in GenBank entry L38423.

A soluble mLT- $\beta$ -R (hIgG1) fusion protein was prepared by PCR amplification of the full length mLT- $\beta$ -R cDNA clone as a template and the primers 5'AACTGCAGCGGCCGCCATGCGCCTGCCC 3' and 5'GACTTTGTCTGACCATTTGCTCCTGGCTCTGGGGG 3'. The amplified product was purified and cut with NotI and SalI and ligated with a SalI/NotI human IgG1 Fc fragment into NotI-linearized and

- 43 -

phosphatase-treated SAB132 to form JLB 122. For stable expression, the NotI cassette containing the mLT- $\beta$ -R-Fc fragment was transferred into the NotI site of pMDR901 forming PSH001 and the vector was transfected into CHO cells as described (Browning et al., J. Immunol., 154, pp. 33-46 (1995)). Cell clones  
5 secreting mLT- $\beta$ -R-Fc were identified by ELISA analysis. The purified receptor fusion protein was isolated from CHO cell supernatants by Protein A Sepharose Fast Flow chromatography (Pharmacia).

10

**EXAMPLE 3****USE OF SOLUBLE HUMAN LT- $\beta$ -R-Fc TO BLOCK  
LT- $\beta$  RECEPTOR-LIGAND INTERACTIONS**

Soluble hLT- $\beta$ -R-Fc was tested for its ability to block  
15 LT ligand binding to the LT- $\beta$  receptor in the tumor cell cytotoxicity assay described above. In this assay, a soluble form of the LT ligand (hLT- $\alpha$ 1/ $\beta$ 2), which activates LT- $\beta$ -R signalling, is used to kill human tumor cells. Inhibitors of LT- $\beta$ -R signalling can reduce LT- $\beta$ -R-induced tumor cell  
20 cytotoxicity.

Soluble LT- $\alpha$ 1/ $\beta$ 2 ligands comprise truncated or modified LT- $\beta$  subunits lacking a functional transmembrane domain. Soluble LT- $\alpha$ 1/ $\beta$ 2 ligands bind to and stimulate LT- $\beta$ -R signalling as well as surface forms of LT ligand (Browning et  
25 al., J. Immunol., 154, pp. 33-46 (1995)).

Serial dilutions of hLT- $\alpha$ 1/ $\beta$ 2, hTNF or hLT- $\alpha$  were prepared in 0.05 ml in 96 well plates and 5000 trypsinized HT29 cells (ATCC) added in 0.05 ml media containing 80 U/ml (antiviral units) of hu-IFN- $\gamma$ . After 4 days, mitochondrial  
30 reduction of the dye MTT was measured as follows: 10  $\mu$ l of MTT was added and after 3 hours, the reduced dye dissolved with 0.09 ml of isopropanol with 10mM HCl, and the O.D. measured at 550 nm. Soluble receptor forms or pure human IgG were added in 10  $\mu$ l prior to the addition of the cells to give a final  
35 concentration of 5  $\mu$ g/ml.

- 44 -

Table 1 compares the ability of hLT- $\beta$ -R-Fc and p55-TNF-R-Fc chimeras (with human IgG as a control) to block the inhibitory effects of various soluble TNF and LT ligands on HT29 tumor cell growth.

5 **Table I**

Ability of LT- $\beta$ -R and p55-TNF-R Immunoglobulin Fusion Proteins to Block the Inhibitory Effects of Various TNF and LT Ligands on HT29 Growth

	Concentration of Cytotoxic Agent (ng/ml) Resulting in 50% Growth Inhibition		
	In the Presence of <sup>a</sup>		
	<u>Cytotoxic Agent</u>	<u>hu-IgG control</u>	<u>p55-TNF-R-Fc</u> <u>LT-<math>\beta</math>-R-Fc</u>
	TNF	0.08	>10 <sup>b</sup> 0.08
	LT- $\alpha$	3	>1000 3
15	LT- $\alpha$ 1/ $\beta$ 2	5	5 >200

<sup>a</sup>Each cytotoxic agent was pre-mixed with the Ig fusion proteins for 10 minutes prior to addition to the cells. The final concentration of fusion protein was 5  $\mu$ g/ml.

20 <sup>b</sup>Higher concentrations were not tested.

The data in Table 1 indicate that the soluble human LT- $\beta$ -R fusion protein (hLT- $\beta$ -R-Fc) can effectively block the interaction between LT ligand (LT- $\alpha$ 1/ $\beta$ 2) and cell surface LT- $\beta$  receptors and is thus a LT- $\beta$ -R blocking agent according to this invention.

25 As expected, the soluble TNF-R fusion protein (p55-TNF-R-Fc) completely blocked TNF-induced growth inhibition by binding to TNF and preventing its interaction with surface TNF receptors. This soluble TNF receptor had no effect on LT ligand-mediated anti-proliferative effects. In contrast, the LT- $\beta$ -R-Fc blocked LT  
30 ligand-induced cytotoxic effects but not those of TNF or LT- $\alpha$ . Thus soluble human LT- $\beta$ -R fusion proteins do not interfere with TNF-R activation by TNF and LT- $\alpha$  ligands.

- 45 -

**EXAMPLE 4****USE OF SOLUBLE MURINE LT- $\beta$ -R-Fc TO BLOCK  
MOUSE LT- $\beta$  RECEPTOR-LIGAND INTERACTIONS**

A soluble murine LT- $\beta$  receptor coupled to a human IgG1 Fc domain (mLT- $\beta$ -R-Fc; see Example 2) was tested for its ability to block LT- $\beta$  receptor-ligand interaction in mouse using a cytotoxicity assay on mouse cells (**Figure 2**). The cytotoxicity assay was performed on WEHI 164 cells using essentially the same procedure as was used in the HT29 cell assay described in Example 3 (see also Browning and Ribolini, J. Immunol., 143, pp. 1859-67 (1989)).

**Figure 2** shows the effects of mLT- $\beta$ -R-Fc on ligand-induced LT- $\beta$ -R signalling in mouse WEHI 164 cells. As this assay indicates, WEHI 164 cells are killed by treating them with LT- $\alpha/\beta$  ligand at concentrations ranging from about 1 to 100 ng/ml. Soluble mLT- $\beta$ -R-Fc (10  $\mu$ /ml) blocks the LT ligand-activated cell death. Adding a soluble mouse p55-TNF-R-Fc fusion protein or IgG control antibodies (each at 10 $\mu$ /ml) had little or no effect on blocking cell death. These data show that the mLT- $\beta$ -R-Fc fusion protein can effectively compete with surface LT- $\beta$ -R molecules for LT ligand binding. These data also show that LT- $\alpha/\beta$ -induced cytotoxicity is LT- $\beta$ -R-mediated and can be inhibited by soluble mLT- $\beta$ -R-Fc, which acts as a LT- $\beta$ -R blocking agent according to the present invention.

25

**EXAMPLE 5****USE OF ANTI-HUMAN LT- $\beta$ -R ANTIBODIES TO BLOCK  
LT- $\beta$  RECEPTOR-LIGAND INTERACTIONS**

Mouse monoclonal antibodies (mAbs) directed against the human LT- $\beta$  receptor were prepared by intraperitoneal immunization of RBF mice repetitively with a CHO cell-derived hLT- $\beta$ -R-Fc fusion protein attached to Protein A Sepharose beads in the absence of adjuvant. Animals were finally boosted with soluble hLT- $\beta$ -R-Fc, both i.p. and i.v., spleen cells were fused via classical protocols and hybridoma supernatants were screened by ELISA (Ling et al., J. Interferon and Cytokine Res., 15, pp. 53-59 (1995)). Hybridoma supernatants were screened further for their ability to block binding of activated II-23 hybridoma

- 46 -

cells -- which express surface LT- $\alpha$ 1/ $\beta$ 2 -- to LT- $\beta$ -R-Fc coated plates in a cell panning assay. Pure mAbs were prepared by Protein A Sepharose purification (Pharmacia) of IgG from culture supernatants.

5 To determine whether an anti-LT- $\beta$  receptor mAb could block LT- $\beta$ -R signalling initiated by the binding of soluble LT, a tumor cell cytotoxicity assay was performed using WiDr human carcinoma cells. In the cytotoxicity assays, serial dilutions of LT- $\alpha$ 1/ $\beta$ 2 were prepared in 0.05 ml in 96 well plates and 10  $\mu$ l  
10 of a 100  $\mu$ g/ml solution containing either control mouse IgG1 mAb or the anti-LT- $\beta$  receptor mAb was added. 5000 trypsinized WiDr cells (ATCC) were then added to each well in 0.05 ml of media containing 50 U/ml (antiviral units) of hu-IFN- $\gamma$ . After 4 days, mitochondrial reduction of the dye MTT was measured as follows:  
15 10  $\mu$ l of MTT was added and after 3 hours, the reduced dye dissolved with 0.09 ml of isopropanol with 10mM HCl, and the O.D. measured at 550 nm. The amount of purple color is proportional to the amount of cell growth.

**Figure 3** shows that the anti-LT- $\beta$ -R mAb BDA8 acts as a  
20 LT- $\beta$ -R blocking agent according to this invention. Human WiDr carcinoma cells stop growing in the presence of IFN- $\gamma$  and soluble LT- $\alpha$ 1/ $\beta$ 2 ligand (from about 0.05 to 50 ng/ml). An IgG1 control antibody (10  $\mu$ g/ml) has no effect on this growth inhibition. In contrast, the anti-LT- $\beta$ -R mAb BDA8 (10  $\mu$ g/ml)  
25 restores the ability of WiDr cells to grow in the presence of soluble LT- $\alpha$ 1/ $\beta$ 2 ligand.

#### EXAMPLE 6

##### USE OF ANTI-HUMAN LT- $\beta$ ANTIBODIES TO BLOCK RECEPTOR-LIGAND INTERACTIONS

30 Anti-human LT- $\beta$  mAbs were prepared by immunizing RBF mice with washed protein A Sepharose-9E10-rLT- $\beta$  beads containing about 1-2  $\mu$ g of human recombinant LT- $\beta$  in CFA, and followed with one boost of the same material in IFA. Eight weeks after the last boost, mice were given i.v. 30  $\mu$ g of purified soluble rLT- $\beta$   
35 (acid eluted off the 9E10 resin) and 20  $\mu$ g of the same soluble material 2 days later. One day after the second i.v. boost, the spleen cells were fused using classical protocols to create



- 47 -

mAbs. Hybridoma supernatants were screened directly by ELISA or by FACS staining of PMA-activated II-23 cells. Pure mAbs were prepared by Protein A Sepharose Fast Flow purification of IgG from culture supernatants (Pharmacia).

5 A FACS assay was used to select antibodies directed against LT- $\beta$  that can effectively block the binding of soluble LT- $\alpha/\beta$  ligand to LT- $\beta$  receptors on the surface of a cell -- thus mimicking the interaction between two cells *in vivo*. In this assay, soluble human LT- $\beta$ -R-Fc (2  $\mu$ g/ml) was allowed to bind to  
10 surface LT ligand on PMA-activated II-23 cells (Browning et al., *J. Immunol.*, 154, pp. 33-46 (1995)) in the presence of increasing concentrations of the test anti-LT- $\beta$  mAb (0.02-20  $\mu$ g/ml). The cells were washed and the bound LT- $\beta$ -R-Fc was detected by reaction with phycoerythrin-labelled donkey anti-  
15 human IgG. The amount of bound fluorescent label was determined by FACS analysis and the mean fluorescence intensity was plotted.

**Figure 4** shows the results of a FACS assay which measured the ability of the anti-LT- $\beta$  mAb B9 to block LT- $\beta$   
20 receptor-ligand interaction as described above. This experiment shows that the anti-LT- $\beta$  mAb B9 (0.02-5  $\mu$ g/ml) can specifically and effectively compete for cell surface LT ligand binding with soluble LT- $\beta$ -R fusion protein (2  $\mu$ g/ml) and thus qualifies as an LT- $\beta$ -R blocking agent according to this invention.

25

#### EXAMPLE 7

##### USE OF ANTI-MOUSE LT- $\alpha/\beta$ ANTIBODIES TO BLOCK RECEPTOR-LIGAND INTERACTIONS

Soluble mouse LT- $\alpha/\hat{a}$  complexes were prepared as described above for the human soluble LT- $\alpha/\beta$  complexes. The  
30 soluble mouse LT- $\hat{a}$  subunit was made based on sequence information previously described (Lawton et al., *J. Immunol.*, 154, pp. 239-46 (1995)). Soluble murine LT- $\alpha/\hat{a}$  complexes were expressed using the baculovirus/insect cell expression system and the LT- $\alpha/\hat{a}$  complexes were isolated by affinity  
35 chromatography using human p55 TNF-R and LT- $\hat{a}$ -R columns essentially as described above for the expression and purification of human LT- $\alpha/\hat{a}$  complexes. Armenian hamsters

- 48 -

were immunized with purified soluble murine LT- $\alpha$ / $\beta$  complex essentially as described in Example 6. Hamster spleen cells were fused to the mouse P3X hybridoma cell line as described (Sanchez-Madrid et al., Methods of Enzymology, 121, pp. 239-44 (1986)). Hybridomas were grouped as anti-mLT- $\hat{\alpha}$  or anti-mLT- $\alpha$  on the basis of their binding characteristics to either the LT- $\alpha$ / $\hat{\alpha}$  complex or to LT- $\alpha$  alone, respectively. Hybridoma cells were expanded and the antibodies purified from the culture supernatant using Protein A affinity chromatography (Pharmacia).

10 To assess whether hamster anti-mouse LT- $\alpha$  and LT- $\beta$  mAbs could block LT ligand binding to mLT- $\hat{\alpha}$ -R, we used TIMI-4 cells (ATCC), a murine T cell line that expresses surface LT ligand following PMA activation for 7 hours. Hamster anti-mLT- $\alpha$  or anti-mLT- $\beta$  mAbs

15 were preincubated with the cells for 30 minutes at 4 C and then washed twice. The washed cells were incubated with 1  $\mu$ g/ml of mLT- $\beta$ -R-Fc at 4 C. After 30 minutes, cells were washed free of unbound mLT- $\beta$ -R-Fc and then incubated for 30 minutes with 10  $\mu$ g/ml of phycoerythrin-labelled donkey anti-human IgG to detect

20 bound mLT- $\beta$ -R-Fc. The amount of bound fluorescent label was determined by FACS analysis and the mean fluorescence intensity was calculated.

Using this analysis, it was found that the hamster anti-mLT- $\hat{\alpha}$  mAb could effectively block soluble LT- $\beta$  receptor

25 binding to T cell surface LT ligand. The results are shown in Table 2.

- 49 -

**Table 2**

Ability of Anti-mouse LT- $\beta$  Monoclonal Antibody To Inhibit mLT- $\beta$ -R-Fc Binding To Murine Surface LT Ligand

Conc. mAb ( $\mu$ g/ml)	Anti-mLT- $\hat{\alpha}$ (BB.F6)		Anti-mLT- $\alpha$ (AF.B3)	
	MFCI <sup>b</sup>	% Inh <sup>c</sup>	MFCI <sup>b</sup>	% Inh <sup>c</sup>
0 <sup>a</sup>	6	-	6	-
0	85	0	85	0
10 0.01	71	18	84	2
0.03	67	23	86	0
0.1	51	44	86	0
0.3	36	62	84	2
1.0	29	71	89	0
15 3.0	17	86	88	0
10.0	11	94	95	0
30.0	10	95	94	0
100.0	8	98	92	0

20 <sup>a</sup>no receptor added

<sup>b</sup>Mean Fluorescence Channel No.

<sup>c</sup>Percent Inhibition

**EXAMPLE 8**

**LT- $\beta$ -R Blocking Agents Inhibit  
Th1-Mediated Contact Hypersensitivity in Mouse**

25 20 g female Balb/c mice (Jackson Laboratories, Bar Harbor, ME) were initially sensitized by applying 25  $\mu$ L of 0.5% 2,4-dinitrofluorobenzene (DNFB) in 4:1 v/v acetone:olive oil onto the bottom of each hind foot. Twenty-four hours after the  
30 initial sensitization, we again sensitized each mouse with 25  $\mu$ L of the same solution. Sensitizations were performed while restraining the unanesthetized mouse. On day 5 (120 hours after the initial sensitization), we anesthetized the mice with  
90:10 mg/kg ketamine:xylazine (i.p.) and applied a sub-irritant  
35 dose of 10  $\mu$ L of 0.2% DNFB to the dorsal and ventral surfaces of the left ear. The right ear received a similar application of the 4:1 v/v acetone:olive oil vehicle.

Four hours after challenging the immune response, we administered increasing concentrations of the mLT- $\beta$ -R-Fc (0.08-  
40 5.0 mg/kg; Example 2) to the mice in 0.1 ml of phosphate

- 50 -

buffered saline (PBS) by intravenous injection. Injections of PBS buffer alone, or 20 mg/kg of a human IgG fusion protein (LFA3-Fc) (Miller et al., J. Exp. Med., 178, pp. 211-22 (1993)) served as negative controls. Injection of 8 mg/kg of an anti-VLA4-specific mAb (PS/2 mAb; Chisolm et al., Eur. J. Immunol., 23, pp. 682-88 (1993)) -- which is known to inhibit CHS by blocking the influx of T cells into the challenge site -- served as a positive control. Groups of four to eight mice were treated per concentration of antibody.

Twenty four hours after challenge, mice were again anesthetized with ketamine:xylazine and the ear thickness of both ears measured with an engineer's micrometer to an accuracy of  $10^{-4}$  inches. The ear swelling response for each mouse was the difference between its control- and DNFB-challenged ear thickness. Typical uninhibited ear swelling responses were  $95-110 \times 10^{-4}$  inches. Inhibition of the ear swelling response was judged by comparison of treated groups with their negative control group. Statistical significance of the difference among treatment groups was evaluated using one-way analysis of variance followed by computation of the Tukey-Kramer Honestly Significant Difference (JMP, SAS Institute) using  $p < 0.05$ .

Figure 5 shows that administering increasing concentrations of mLT- $\beta$ -R-Fc causes a significant reduction in the ear swelling response of DNFB-treated mice compared to uninhibited DNFB-treated control animals (PBS and LFA3-Fc). Soluble LT- $\beta$ -R (from about 1-5 mg/kg) can block this contact DTH reaction as effectively as the inhibitor anti-VLA4-specific mAb. The portion of this ear swelling assay which is not inhibited probably results from "nonspecific" granulocyte infiltration.

#### Example 11

##### Dextran Sulphate Solution (DSS) IBD Model

Mice were treated as defined in the figure legend with hLFA3-Ig, i.e. a control Ig fusion protein or mLT $\beta$ R-Ig by intraperitoneal injection. At day 0, the drinking water was changed to a 5% dextran sulphate solution and the mice were left

- 51 -

on this fluid for one week. One week later, i.e. 2 weeks after starting DSS administration, mice were sacrificed and the weight change and the large bowel length (from anus to cecum) measured. Figure \_\_ shows the weight changes and bowel lengths after various treatments. The shortened bowel length as well as the weight loss is indicative of IBD. It was seen that mLT $\beta$ R-Ig treatment dramatically prevents the colon shortening and weight loss indicating efficacy.

Figure 6:

The weight change observed 14 days post initiation of DSS in the drinking water following various treatments. Veh = vehicle, LT $\beta$ R and LFA3 refer to mLT $\beta$ R-Ig and hLFA3-Ig fusion proteins that were administered by intraperitoneal injection of 100 ug 1 week prior to adding DSS, at the point of DSS administration and 1 week later (i.e. 3 injections at -1, 0 and 1 week). There were 10 animals per group.

Figure 7:

The colon length at 14 days post DSS administration following the various treatments described in 6.

20

#### **Example 10** **CD45RB<sup>hi</sup>/scid model of IBD**

CD4 positive T cells are isolated from C.B-17 female mice using magnetic bead technology as described earlier (F. Powrie et al International Immunology 5:1461-1471 1993). The CD4 cells depleted of CD8 positive T cells, B cells and monocytes were then sorted by fluorescence activated cell sorting into CD45RB<sup>high</sup> and CD45RB<sup>low</sup> populations also essentially as described above. 5 x 10<sup>5</sup> CD45RB cells were injected intravenously into female C.B-17 scid mice and the body weight of the mice was followed. It can be seen that animals reconstituted with CD45RB<sup>low</sup> cells gained weight in a normal manner. In contrast, animals receiving CD45RB<sup>high</sup> cells eventually lost weight and at 10 weeks were near death. When the control mice had lost roughly 20% of their starting weight, the mice were sacrificed and various organs

- 52 -

analyzed by histology. Typically diseased animals looked cachectic, had diarrhea and had dramatically enlarged colons and ceca. Animals treated as described in the figure legend with hLFA3-Ig were similar to untreated animals, whereas animal  
5 treated with mLT $\beta$ R-Ig had not undergone weight loss, had relatively normal sized colons and lacked the massive inflammatory infiltrates typically observed in the colon. Figure 8 shows the time course of weight loss in CD45RB<sup>high</sup> injected animals treated in various ways and figure 9 shows the  
10 final body weights at 8 weeks post injection. The efficacy of mLT $\beta$ R-Ig in two very different models of IBD, i.e. the CD45RB and DSS models, represents strong evidence for a profound effect of this treatment on the immune system.

Figure 8:

15 Time course of the body weigh following injection of CD45RB CD4 positive T cells into scid mice. Each curve represents one animal and the inscriptions in the panels refer to which cells were injected i.e. CD45RB<sup>high or low</sup> and the nature of the treatment. Animal were treated weekly with 100 ug of protein  
20 injected intraperitoneally. Treatment started 2 weeks prior to the injection of the cells and continued throughout the course of the experiment.

Figure 9:

Mean and standard deviations of the body weights observed  
25 following various treatments at 10 weeks post transplantation (5-6 animals per group).

#### Example 11:

##### A SRBC Model of Delayed Type Hypersensitivity

Female balb/c mice are sensitized by subcutaneous injection  
30 of  $2 \times 10^7$  washed sheep red blood cells (SRBC) in PBS. After 5 days, mice are challenged with a injection of  $1 \times 10^8$  SRBC in PBS into the right footpad (subplantar injection). Various times after injection into the footpad, the footpad thickness was measured with calipers. Figure 10 shows the footpad  
35 swelling response in mice either treated by intraperitoneal

- 53 -

injection with mLT $\beta$ R-Ig. Treatment with mLT $\beta$ R-Ig either at the point of sensitization or at both sensitization or challenge stages inhibited the SRBC induced DTH response.

Figure 10:

- 5 Shown is the increase in footpad thickness measured 18 h post injection with SRBC challenge. Treatments were either a negative control injection of PBS, a positive control antibody PS/2 that blocks VLA4 interactions and hence cell trafficking and mLT $\beta$ R-Ig (100 ug intravenous injections) given either immediately prior  
10 to the sensitizing subcutaneous injection of SRBC, at the challenge point or at both times.

- 54 -

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

- (i) APPLICANT: BROWNING, Jeffrey L.  
BENJAMIN, Christopher D.  
HOCHMAN, Paula S.
- (ii) TITLE OF INVENTION: SOLUBLE LYMPHOTOXIN-BETA RECEPTORS AND  
ANTI-LYMPHOTOXIN RECEPTOR AND LIGAND ANTIBODIES AS  
THERAPEUTIC AGENTS FOR THE TREATMENT OF IMMUNOLOGICAL  
DISEASE
- (iii) NUMBER OF SEQUENCES: 1
- (iv) CORRESPONDENCE ADDRESS:
  - (A) ADDRESSEE: James F. Haley, Jr.
  - (B) STREET: 1251 Avenue of the Americas
  - (C) CITY: New York
  - (D) STATE: New York
  - (E) COUNTRY: U.S.A.
  - (F) ZIP: 10020
- (v) COMPUTER READABLE FORM:
  - (A) MEDIUM TYPE: Floppy disk
  - (B) COMPUTER: IBM PC compatible
  - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
  - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
  - (A) APPLICATION NUMBER:
  - (B) FILING DATE: CONCURRENTLY HERewith
  - (C) CLASSIFICATION:
- (vii) PRIOR APPLICATION DATA:
  - (A) APPLICATION NUMBER: US 08/505,606
  - (B) FILING DATE: 21-JUL-1995
- (viii) ATTORNEY/AGENT INFORMATION:
  - (A) NAME: HALEY, Jr., James F.
  - (B) REGISTRATION NUMBER: 27,794
  - (C) REFERENCE/DOCKET NUMBER: B191CIP PCT
- (ix) TELECOMMUNICATION INFORMATION:



- 55 -

- (A) TELEPHONE: (212) 596-9000  
 (B) TELEFAX: (212) 596-9090  
 (C) TELEX: 14-8367

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 197 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS:  
 (D) TOPOLOGY: linear

## (ii) MOLECULE TYPE: peptide

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Ser	Gln	Pro	Gln	Ala	Val	Pro	Pro	Tyr	Ala	Ser	Glu	Asn	Gln	Thr	Cys	1	5	10	15
Arg	Asp	Gln	Glu	Lys	Glu	Tyr	Tyr	Glu	Pro	Gln	His	Arg	Ile	Cys	Cys	20	25	30	
Ser	Arg	Cys	Pro	Pro	Gly	Thr	Tyr	Val	Ser	Ala	Lys	Cys	Ser	Arg	Ile	35	40	45	
Arg	Asp	Thr	Val	Cys	Ala	Thr	Cys	Ala	Glu	Asn	Ser	Tyr	Asn	Glu	His	50	55	60	
Trp	Asn	Tyr	Leu	Thr	Ile	Cys	Gln	Leu	Cys	Arg	Pro	Cys	Asp	Pro	Val	65	70	75	80
Met	Gly	Leu	Glu	Glu	Ile	Ala	Pro	Cys	Thr	Ser	Lys	Arg	Lys	Thr	Gln	85	90	95	
Cys	Arg	Cys	Gln	Pro	Gly	Met	Phe	Cys	Ala	Ala	Trp	Ala	Leu	Glu	Cys	100	105	110	
Thr	His	Cys	Glu	Leu	Leu	Ser	Asp	Cys	Pro	Pro	Gly	Thr	Glu	Ala	Glu	115	120	125	

- 56 -

Leu Lys Asp Glu Val Gly Lys Gly Asn Asn His Cys Val Pro Cys Lys  
130 135 140

Ala Gly His Phe Gln Asn Thr Ser Ser Pro Ser Ala Arg Cys Gln Pro  
145 150 155 160

His Thr Arg Cys Glu Asn Gln Gly Leu Val Glu Ala Ala Pro Gly Thr  
165 170 175

Ala Gln Ser Asp Thr Thr Cys Lys Asn Pro Leu Glu Pro Leu Pro Pro  
180 185 190

Glu Met Ser Gly Thr  
195

- 57 -

**CLAIMS**

What is claimed is:

- 1        1.    A method for treating or reducing the advancement,  
2    severity or effects of an immunological disease in an animal  
3    comprising the step of administering a pharmaceutical composition  
4    which comprises a therapeutically effective amount of a LT- $\beta$ -R  
5    blocking agent and a pharmaceutically acceptable carrier.
- 1        2.    The method according to claim 1, wherein the LT- $\beta$ -R  
2    blocking agent is selected from the group consisting of a soluble  
3    lymphotoxin- $\beta$  receptor, an antibody directed against LT- $\beta$  receptor,  
4    and an antibody directed against surface LT ligand.
- 1        3.    The method according to claim 2, wherein the animal is  
2    a mammal.
- 1        4.    The method according to claim 3, wherein the mammal is  
2    a human.
- 1        5.    The method according to claim 1, wherein the LT- $\beta$ -R  
2    blocking agent comprises a soluble lymphotoxin- $\beta$  receptor having  
3    a ligand binding domain that can selectively bind to a surface LT  
4    ligand.
- 1        6.    The method according to claim 5, wherein the soluble  
2    lymphotoxin- $\beta$  receptor further comprises a human immunoglobulin Fc  
3    domain.
- 1        7.    The method according to claim 1, wherein the LT- $\beta$ -R  
2    blocking agent comprises a monoclonal antibody directed against LT-  
3     $\beta$  receptor.
- 1        8.    The method according to claim 7, wherein the composition  
2    is administered in an amount sufficient to coat LT- $\beta$  receptor-  
3    positive cells for 1 to 14 days.

- 58 -

1        9. The method according to claim 4, wherein the LT- $\beta$ -R  
2 blocking agent comprises anti-human LT- $\beta$ -R mAb BDA8.

1        10. The method according to claim 1, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against  
3 surface LT ligand.

1        11. The method according to claim 10, wherein the composition  
2 is administered in an amount sufficient to coat surface LT ligand-  
3 positive cells for 1 to 14 days.

1        12. The method according to claim 10, wherein the antibody  
2 is directed against a subunit of the LT ligand.

1        13. The method according to claim 4, wherein the LT- $\beta$ -R  
2 blocking agent comprises anti-human LT- $\beta$  mAb B9.

1        14. The method according to claim 3, wherein the mammal is  
2 a mouse and the LT- $\beta$ -R blocking agent comprises a monoclonal  
3 antibody directed against a murine surface LT ligand.

1        15. A method for inhibiting a Th1 cell-mediated immune  
2 response in an animal comprising the step of administering a  
3 pharmaceutical composition which comprises an effective amount of  
4 a LT- $\beta$ -R blocking agent and a pharmaceutically effective carrier.

1        16. The method according to claim 15, wherein the LT- $\beta$ -R  
2 blocking agent is selected from the group consisting of a soluble  
3 lymphotoxin- $\beta$  receptor, an antibody directed against LT- $\beta$  receptor,  
4 and an antibody directed against surface LT ligand.

1        17. The method according to claim 16, wherein the animal is  
2 a mammal.

1        18. The method according to claim 17, wherein the mammal is  
2 a human.

- 59 -

1        19. The method according to claim 15, wherein the LT- $\beta$ -R  
2 blocking agent comprises a soluble lymphotoxin- $\beta$  receptor having  
3 a ligand binding domain that can selectively bind to a surface LT  
4 ligand.

1        20. The method according to claim 19, wherein the soluble  
2 lymphotoxin- $\beta$  receptor further comprises a human immunoglobulin Fc  
3 domain.

1        21. The method according to claim 15, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against LT-  
3  $\beta$  receptor.

1        22. The method according to claim 21, wherein the composition  
2 is administered in an amount sufficient to coat LT- $\beta$  receptor-  
3 positive cells for 1 to 14 days.

1        23. The method according to claim 18, wherein the LT- $\beta$ -R  
2 blocking agent comprises anti-human LT- $\beta$ -R mAb BDA8.

1        24. The method according to claim 15, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against  
3 surface LT ligand.

1        25. The method according to claim 24, wherein the composition  
2 is administered in an amount sufficient to coat surface LT ligand-  
3 positive cells for 1 to 14 days.

4        26. The method according to claim 24, wherein the antibody  
5 is directed against a subunit of the LT ligand.

1        27. The method according to claim 18, wherein the LT- $\beta$ -R  
2 blocking agent comprises anti-human LT- $\beta$  mAb B9.

1        28. The method according to claim 17, wherein the mammal is  
2 a mouse and the LT- $\beta$ -R blocking agent comprises a monoclonal  
3 antibody directed against a murine surface LT ligand.

- 60 -

1        29. The method according to claim 15, wherein the Th1 cell-  
2 mediated immune response contributes to a delayed type  
3 hypersensitivity reaction.

1        30. The method according to claim 29, wherein the delayed  
2 type hypersensitivity reaction is contact hypersensitivity.

1        31. The method according to claim 29, wherein the delayed  
2 type hypersensitivity reaction is tuberculin-type hypersensitivity.

1        32. The method according to claim 29, wherein the delayed  
2 type hypersensitivity reaction is a granulomatous reaction.

1        33. The method according to claim 15, wherein the Th1 cell-  
2 mediated immune response contributes to cellular rejection of  
3 tissue in the animal after the animal receives a tissue graft.

1        34. The method according to claim 15, wherein the Th1 cell-  
2 mediated immune response contributes to organ rejection in the  
3 animal after the animal receives an organ transplant.

1        35. The method according to claim 15, wherein the Th1 cell-  
2 mediated immune response contributes to an autoimmune disorder in  
3 the animal.

1        36. The method according to claim 35, wherein the autoimmune  
2 disorder is selected from the group consisting of multiple  
3 sclerosis, insulin-dependent diabetes, sympathetic ophthalmia,  
4 uveitis and psoriasis.

1        37. The method according to claim 15, wherein the Th1 cell-  
2 mediated immune response is inhibited without inhibiting a Th2  
3 cell-dependent immune response.

1        38. A pharmaceutical composition comprising a therapeutically  
2 effective amount of a LT- $\beta$ -R blocking agent and a pharmaceutically  
3 acceptable carrier.

- 61 -

1       39. The composition according to claim 38, wherein the LT- $\beta$ -R  
2 blocking agent is selected from the group consisting of a soluble  
3 lymphotoxin- $\beta$  receptor, an antibody directed against LT- $\beta$  receptor,  
4 and an antibody directed against surface LT ligand.

1       40. The composition according to claim 38, wherein the  
2 soluble lymphotoxin- $\beta$  receptor comprises a LT- $\beta$ -R ligand binding  
3 domain that can selectively bind to a surface LT ligand.

1       41. The composition according to claim 40, wherein the  
2 soluble lymphotoxin- $\beta$  receptor further comprises a human  
3 immunoglobulin Fc domain.

1       42. The composition according to claim 38, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against LT-  
3  $\beta$  receptor.

1       43. The composition according to claim 42, wherein the  
2 monoclonal antibody is anti-human LT- $\beta$ -R mAb BDA8.

1       44. The composition according to claim 38, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against  
3 surface LT ligand.

4       45. The composition according to claim 44, wherein the  
5 antibody is directed against a subunit of the LT ligand.

1       46. The composition according to claim 45, wherein the  
2 monoclonal antibody is anti-human LT- $\beta$  mAb B9.

1       47. The composition according to claim 38, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against a  
3 murine surface LT ligand.

1       48. The composition according to claim 42, wherein the  
2 antibody is present in an amount sufficient to coat LT- $\beta$  receptor-  
3 positive cells for 1 to 14 days.

- 62 -

1        49. The composition according to claim 44, wherein the  
2 antibody is present in an amount sufficient to coat surface LT  
3 ligand-positive cells for 1 to 14 days.

1        50. A method for selecting a LT- $\beta$ -R blocking agent comprising  
2 the steps of:

3            a) culturing tumor cells in the presence of an  
4 effective amount of at least one LT- $\beta$ -R activating agent and a  
5 putative LT- $\beta$ -R blocking agent; and

6            b) determining whether the putative LT- $\beta$ -R blocking  
7 agent decreases the anti-tumor activity of the LT- $\beta$ -R activating  
8 agent.

1        51. The method according to claim 50, wherein the LT- $\beta$ -R  
2 activating agent comprises a LT- $\alpha/\beta$  heteromeric complex.

1        52. The method according to claim 51, wherein the LT- $\alpha/\beta$   
2 heteromeric complex has a LT- $\alpha 1/\beta 2$  stoichiometry.

1        53. The method according to claim 50, wherein the LT- $\beta$ -R  
2 activating agent comprises an anti-LT- $\beta$ -R antibody that stimulates  
3 LT- $\beta$ -R signalling.

1        54. A method for inhibiting LT- $\beta$ -R signalling without  
2 inhibiting TNF-R signalling comprising the step of administering  
3 to a subject an effective amount of a LT- $\beta$ -R blocking agent.

1        55. The method according to claim 54, wherein the LT- $\beta$ -R  
2 blocking agent is selected from the group consisting of a soluble  
3 lymphotoxin- $\beta$  receptor, an antibody directed against LT- $\beta$  receptor,  
4 and an antibody directed against surface LT ligand.

1        56. The method according to claim 54, wherein the subject  
2 comprises one or more cells from a mammal.



- 63 -

1        57. The method according to claim 56, wherein the mammal is  
2 a human.

1        58. The method according to claim 54, wherein the LT- $\beta$ -R  
2 blocking agent comprises a soluble lymphotoxin- $\beta$  receptor having  
3 a ligand binding domain that can selectively bind to a surface LT  
4 ligand.

1        59. The method according to claim 58, wherein the soluble  
2 lymphotoxin- $\beta$  receptor further comprises a human immunoglobulin Fc  
3 domain.

1        60. The method according to claim 54, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against LT-  
3  $\beta$  receptor.

1        61. The method according to claim 57, wherein the LT- $\beta$ -R  
2 blocking agent comprises anti-human LT- $\beta$ -R mAb BDA8.

1        62. The method according to claim 54, wherein the LT- $\beta$ -R  
2 blocking agent comprises a monoclonal antibody directed against  
3 surface LT ligand.

1        63. The method according to claim 62, wherein the antibody  
2 is directed against a subunit of the LT ligand.

1        64. The method according to claim 57, wherein the LT- $\beta$ -R  
2 blocking agent comprises anti-human LT- $\beta$  mAb B9.

1        65. The method according to claim 56, wherein the mammal is  
2 a mouse and the LT- $\beta$ -R blocking agent comprises a monoclonal  
3 antibody directed against a murine surface LT ligand.

1        66. The method according to claims 60, wherein the LT- $\beta$ -R  
2 blocking agent is administered in an amount sufficient to coat LT- $\beta$   
3 receptor-positive cells for 1 to 14 days.

- 64 -

1        67. The method according to claims 62, wherein the LT- $\beta$ -R  
2 blocking agent is administered in an amount sufficient to coat  
3 surface LT ligand-positive cells for 1 to 14 days.

1        68. A method of treating inflammatory bowel syndrome  
2 comprising administering a therapeutically effective amount of an  
3 LT- $\beta$ -R fusion protein.

1        69. The method of claim 68 wherein the fusion protein is LT-  
2  $\beta$ -R a fusion of and an immunoglobulin Fc domain.

1	SQPQAVPPYA	SENQTCRDQE	KEYYEPQHRI	CCSRCPPGTY	VSAKCSRIRD	50
51	TVCATCAENS	YNEHWNYLTI	CQLCRPCDPV	MGLEEIAPCT	SKRKTQCRCQ	100
101	PGMFCAAWAL	ECTHCELLSD	CPPGTEAELK	DEVGKGNNHC	VPCKAGHFQN	150
151	TSSPSARCQP	HTRCENQGLV	EAAPGTAQSD	TTCKNPLEPL	PPEMSGT	197

FIG. 1




FIG. 2

2/6




FIG. 3




FIG. 4

3/6




FIG. 5




FIG. 6

4/6




FIG. 7

5/6

### Protective Effect of mLT $\beta$ R-Ig in the CD45RB high Murine Model of IBD






FIG. 8

6/6

EFFECT OF mLTBR-Ig ON MOUSE  
BODY WEIGHT AT 10 WEEKS POST  
TRANSPLANTATION OF CD45RB<sup>hi</sup> CELLS





# INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 96/12010

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K38/17 A61K39/395 G01N33/577 G01N33/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>WO 94 13808 A (BIOGEN ET AL.) 23 June 1994 cited in the application</p> <p>see claims 24-27</p> <p style="text-align: center;">--- -/--</p>	<p>1-4, 10-18, 24-39, 44-47, 49, 54-57, 62-65,67</p>

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \* "A" document defining the general state of the art which is not considered to be of particular relevance
- \* "E" earlier document but published on or after the international filing date
- \* "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \* "O" document referring to an oral disclosure, use, exhibition or other means
- \* "P" document published prior to the international filing date but later than the priority date claimed

- \* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* "&" document member of the same patent family

Date of the actual completion of the international search

14 November 1996

Date of mailing of the international search report

29. 11. 96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax (+ 31-70) 340-3016

Authorized officer

Nooij, F

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 96/12010

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	THE JOURNAL OF IMMUNOLOGY, vol. 154, no. 1, 1 January 1995, BALTIMORE, MD, USA, pages 33-46, XP002018575 J. BROWNING ET AL.: "Characterization of surface lymphotoxin forms. Use of specific monoclonal antibodies and soluble receptors." cited in the application see the whole document ---	1-6, 10-20, 24-28, 38-41, 44-47, 49-52, 54-59, 62-69
P,X	EUROPEAN CYTOKINE NETWORK, vol. 7, no. 2, April 1996, MONTROUGE, page 167 XP000611207 P. RENNERT ET AL.: "Normal development of lymph nodes is disrupted by soluble LT beta receptor - Ig fusion protein." see abstract 17 ---	1-3,5,6, 15-17, 19,20, 29-41, 54-56, 58,68,69
P,X	THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 183, no. 3, 1 March 1996, NEW YORK, NY, USA, pages 867-878, XP000608398 J. BROWNING ET AL.: "Signaling through the lymphotoxin beta receptor induces the death of some adenocarcinoma tumor lines." see the whole document -----	1-9, 15-23, 29-43, 48-61, 66-69

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/ 12010

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim(s) 1-37 and 54-69  
is(are) directed to a method of treatment of the human/animal  
body, the search has been carried out and based on the alleged  
effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such  
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all  
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment  
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report  
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is  
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/12010

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9413808	23-06-94	AU-A- 6048394	04-07-94
		CA-A- 2150249	23-06-94
		EP-A- 0672143	20-09-95
		JP-T- 8507201	06-08-96
-----			

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**